


Phenolic compounds and multifunctional properties of thinned apples: revalorizing fruit by-products for industrial applications

Javier Cano-Lou, Gema Casado-Hidalgo, Francisco Les, Ana Pina, Laura Acquaticci, Giovanni Caprioli, Víctor López



PII: S2590-1575(26)00161-6

DOI: <https://doi.org/10.1016/j.fochx.2026.103640>

Reference: FOCHX 103640

To appear in: *Food Chemistry: X*

Received date: 5 September 2025

Revised date: 28 January 2026

Accepted date: 3 February 2026

Please cite this article as: J. Cano-Lou, G. Casado-Hidalgo, F. Les, et al., Phenolic compounds and multifunctional properties of thinned apples: revalorizing fruit by-products for industrial applications, *Food Chemistry: X* (2024), <https://doi.org/10.1016/j.fochx.2026.103640>

This is a PDF of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability. This version will undergo additional copyediting, typesetting and review before it is published in its final form. As such, this version is no longer the Accepted Manuscript, but it is not yet the definitive Version of Record; we are providing this early version to give early visibility of the article. Please note that Elsevier's sharing policy for the Published Journal Article applies to this version, see: <https://www.elsevier.com/about/policies-and-standards/sharing#4-published-journal-article>. Please also note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Phenolic compounds and multifunctional properties of thinned apples: revalorizing fruit by-products for industrial applications

Javier Cano-Lou ^{a, b}, Gema Casado-Hidalgo ^a, Francisco Les^{a, b}, Ana Pina ^c, Laura
Acquaticci ^d, Giovanni Caprioli ^d, Víctor López ^{a, b *}

* Corresponding author

^a *Department of Pharmacy, Faculty of Health Sciences, Universidad San Jorge, 50830, Zaragoza, Spain.*

^b *Institute of agri-food research of Aragón, IA2, Zaragoza University-CITA, 50013, Zaragoza, Spain.*

^c *Unidad de Hortofruticultura, Centro de Investigación y Tecnología Agroalimentaria de Aragón (CITA), Av Montañana 930, 50059, Zaragoza, Spain*

^d *Chemistry Interdisciplinary Project (ChIP), School of Pharmacy, University of Camerino, Via Madonna delle Carceri, 62032, Camerino, Italy.*

Abstract

Thinned apples are a common by-product of apple orchards, produced in abundant quantities through thinning or natural drop, leading to the loss of substantial amounts of phenolic compounds with potential bioactive properties. Ten thinned apple samples, representing both commercial and autochthonous cultivars, were ultrasonicated using ethanol as green solvent, with the main goal of reusing their phenolic content for industrial and nutritional applications. For this reason, the phenolic content was characterized with high-performance liquid chromatography coupled with tandem mass spectrometry

(HPLC-MS/MS). The antioxidant, antidiabetic, and cardiovascular activities were evaluated and correlated with the targeted phenolic compounds. Under non-cytotoxic conditions in Caco-2, HepG2 and HFF-1 cells, Amarilla de Octubre and Pomera de Pomes Agrias showed the strongest antioxidant-antiglycation activity as well as ACE inhibition, while Reineta and Verde Doncella excelled in HMGR and ACE inhibition. Pomera de Pomes Agrias, Vadiello-02, Reineta and Verde Doncella led digestive enzymes under Caco-2 safety.

Keywords

Nutraceuticals, by-products, circular economy, polyphenols, ultrasound-assisted extraction, *in vitro* bioactivity, cytotoxicity.

1. Introduction

Metabolic disorders are non-communicable diseases with a great impact nowadays, particularly in developed countries (Noubiap et al., 2022). Between them, metabolic syndrome (MetS) is a complex health disorder characterised by the synergy of dysglycemia, severe overweight, hypertension, hypertriglyceridemia and decreased high-density lipoprotein (HDL) cholesterol levels (Grima Serrano, 2010). Hyperglycaemia and obesity are major contributors to MetS by promoting oxidative stress, which drive to atherogenic dyslipidaemia, insulin resistance and hypertension (Masenga et al., 2023).

The rising prevalence of several of these risk factors have triggered a major scientific interest in functional food compounds. Particularly, regarding the role of plant-derived bioactive compounds, including phytochemicals, organic acids, dietary fiber, amino acids, monosaccharides and polysaccharides (Dou et al., 2025) in the modulation of metabolic pathways associated with the prevention and management of MetS (Capanoglu et al., 2022).

Among these, phenolic compounds, non-nutritive phytochemicals commonly found in plants classified as phenolic acids, flavonoids, tannins, stilbenes coumarins and lignans, are widely recognized for their antioxidant activity and their capacity to diminish oxidative stress and the risk of MetS (Tian et al., 2020). In particular, dietary flavonoid subclasses, including flavones, flavanones, flavanols, flavonols, isoflavones, anthocyanins or anthocyanidins are the most abundant group of polyphenols. These subclasses share a core of diphenylpropane ($C_6-C_3-C_6$) structure formed by a pair of aromatic rings linked by a three-carbon bridge that defines the central C ring (Jomova et al., 2025) and are closely associated with obesity reduction, improved lipid profiles, blood glucose control and blood pressure regulation (Khan et al., 2021). Similarly, phenolic acids, which represent about one-third of dietary phenolics (Yang et al., 2001)

are associated with metabolic health benefits and are mainly classified into benzoic and cinnamic acids, both featuring a benzene ring and a carboxylic group, often with hydroxyl and/or methoxyl substitutions (Nunes et al., 2025).

Phenolic compounds are commonly contained in food of plant origin, but compared to other plant sources, phenolic content is particularly high in unripe fruits, which are often discarded as by-products during the thinning process. While thinning improves crop quality by reallocating resources to the remaining fruits, it tends to overlook the potential value of these phenolic-rich discarded fruits (Wei et al., 2021). Specifically, thinned apples are a considerable portion of apple production representing between 20 and 30 percent and tend to show higher concentrations of total phenolic compounds, flavonoids, and proanthocyanidins, as well as enhanced antioxidant activity, compared to mature apples (Zheng et al., 2012). Indeed, the valorization of this type of fruit by-products depends on the specific properties of the plant matrix and the targeted functional compounds (Ran et al., 2019).

The recovery of phenolic compounds can be achieved through novel approaches such as ultrasound-assisted extraction (UAE) which enhance the penetration of various solvents and provides higher yields along with several procedural advantages (Nirmal et al., 2023). In particular ethanol is a safe and efficient green solvent that typically yields phenolic enriched extracts with higher antioxidant activity, whereas adding water may increase total yield but dilute the phenolic fraction and weaken antioxidant related responses due to co extraction of abundant low activity matrix components that mask phenolic driven effects (Do et al., 2014).

In addition, the phenolic extracts obtained can be processed into dietary supplements as nutraceuticals for the treatment and prevention of MetS (Popiolek-Kalisz & Glibowski,

2023), and although still under investigation, they may offer an accessible, safe and novel therapy (Fahed et al., 2022).

This manuscript aims to evaluate whether thinned apple by-products from both autochthonous and commercial apple cultivars represent valuable natural sources of phenolic bioactive compounds, and to explore their biological properties and specific mechanisms of action in MetS along with evidence of safety in intestinal, hepatic and structural *in vitro* cell models, emphasizing their relevance for industrial applications through a sustainable and circular economy. We hypothesize that a metabolomics-driven profiling strategy, integrating LC–MS/HPLC chemical targeted analysis with chemometric modelling, will robustly link extract composition to pharmacodynamic profile (IC₅₀), enabling early identification of the metabolite features most predictive of activity. Consequently, this framework should support extract prioritization and guide targeted isolation of bioactive constituents thereby accelerating the discovery pipeline. For this purpose, their phenolic characterization was performed, along with the *in vitro* assessment of its antioxidant, antidiabetic, antiglycant, hypolipidemic, hypocholesterolemic, and hypotensive activities.

2. Materials and methods

2.1. Thinned apples samples preparation

Eight distinct cultivars of thinned apples (Reineta, Verde Doncella, Golden Smoothie, Fuji and Royal Gala as commercial cultivars; and Daroca-02, Camuesa de Castellano de Daroca and Vadiello-02 as autochthonous cultivars) were collected from Zaragoza (Aragón, Spain). Nevertheless, two specific varieties of autochthonous thinned apples (Amarilla de Octubre and Pomera de Pomes Agrias) were collected from the Garcipollera Valley, located in Huesca (Aragón, Spain). The thinning period took place in May 2024, and the samples of each variety were collected from two different trees and various parts

of each tree. Subsequently, the thinned apples were frozen at $-80\text{ }^{\circ}\text{C}$ and lyophilized with LyoMicron $-85\text{ }^{\circ}\text{C}$ freeze-drying equipment (Dara-Lyo brand, Dara Pharma Group, Barcelona, Spain) for five days until completely dried. The dried samples were stored at $-40\text{ }^{\circ}\text{C}$ in hermetically sealed jars with silica gel.

2.2. Extraction of phenolic compounds

After a proper drying and storage, the samples were pulverized with a laboratory mill before the extraction method was applied. The procedure used was the ultrasound-assisted extraction (UAE) with the sonicator 505 (Fisherbrand™, Fisher Scientific, Madrid, Spain) equipped with a microtip of 13 mm diameter. The phenolic extraction process was achieved through the mixture of the powdered sample and absolute ethanol as green solvent, maintaining a solid liquid ratio (SLR) of 1:50 along with an exposure time of 20 minutes with an ultrasound amplitude of 40%. For each ratio, 5 g of powder were exactly weight and combined with 250 mL of absolute ethanol. To prevent overheating, the sample mixture was kept cool using ice bath throughout the extraction process. UAE conditions were preselected to ensure standardized extraction across apple cultivar by-products, and their plausibility defined within literature-reported operating ranges for phenolic extraction to balance mass transfer, rapid phenolic release, and moderate cavitation intensity, while limiting over-processing (Medina-Torres et al., 2017).

Every extracted solution mixture was centrifugated for 5 minutes, and the supernatant filtered by vacuum filtration with a $0.45\text{ }\mu\text{m}$ pore-sized polyamide 66 nylon membrane and a thickness of 90–140 μm . Additionally, the absolute ethanol was evaporated by rotary evaporator (Buchi UK Ltd, Newmarket, United Kingdom) until a semi-solid final mixture was obtained from each thinned apple sample. The extracts were stored in hermetic falcon tubes at -20°C . The extraction yield, defined on a dry-weight basis as the mass of dried ethanolic extract recovered relative to the initial mass of lyophilized

thinned-apple material, was approximately 30% (w/w) for most cultivars. In contrast, the autochthonous cultivars Amarilla de Octubre, Pamera de Pomes Agrias and Camuesa de Castellano de Daroca yielded approximately 20% (w/w). To contextualize yield in terms of the target phytochemical fraction, the summed targeted phenolic content quantified by HPLC–MS ranged from 9687.52 to 13167.26 mg kg⁻¹ in the 20% yield extracts, and from 8105.21 to 14010.93 mg kg⁻¹ in the 30% yield extracts. The lower yields observed in some autochthonous cultivars likely reflect differences in matrix properties rather than a reduced phenolic content.

2.3. Analysis of phenolic compounds by HPLC-MS/MS

2.3.1. Reagents and standards

A set of 38 reference compounds was utilized and obtained from PhytoLab (Vestenbergsgreuth, Germany) and Sigma-Aldrich (Milan, Italy). Stock solutions of pure compounds, formulated at 1000 mg L⁻¹ in methanol of HPLC grade, served as the basis for subsequent analysis. These solutions were kept in glass containers with storage conditions at 4 °C for most solutions, while anthocyanins were preserved at -15 °C to prevent degradation. The daily preparation of standard working solutions at various concentrations was carried out by mixing stock solutions with HPLC-grade methanol. Formic acid (99%) was supplied by Merck (Darmstadt, Germany), and hydrochloric acid (37%, analytical grade) was sourced from Carlo Erba Reagents (Milan, Italy). HPLC-grade methanol was acquired from Sigma-Aldrich (Milan, Italy). Deionized water with a resistivity greater than 18 MΩ·cm was processed using a Milli-Q SP Reagent Water System (Millipore, Bedford, MA, USA). Solvents and prepared solutions were passed through 0.2 μm polyamide membrane filters acquired from Sartorius Stedim (Goettingen, Germany). The filtration of the samples was carried out using Phenex™ RC syringeless

filters (4 mm, 0.2 μm) provided by Phenomenex (Castel Maggiore, BO, Italy), prior to the analytical assay.

2.3.2 HPLC-ESI-MS/MS instrument and analysis

The analyse was carried out by an Agilent 1290 Infinity system coupled to a Triple Quadrupole 6420 mass spectrometer (Agilent Technologies, Santa Clara, CA), fitted with an electrospray ionization (ESI) source working in positive and negative ion mode. The method applied was previously developed and validated (Mustafa et al., 2022). Optimization of MS/MS parameters for target compounds was performed via flow injection analysis (FIA), using 1 μL injections of individual standard solutions (10 mg L^{-1}), with the aid of Agilent's Optimizer Software.

The separation of target analytes was carried out using a Synergi Polar-RP C18 analytical column (250 mm \times 4.6 mm, 4 μm ; Phenomenex, Cheshire, UK) and a Polar RP guard cartridge (4 mm \times 3 mm ID) previously assembled. The mobile phase consisted of a solution of (A) water and (B) methanol, containing 0.1% formic acid, delivered at an elution rate of 0.8 mL min^{-1} under gradient conditions. The gradient program was structured as follows: an initial isocratic phase at 20% B from 0 to 1 minute; a linear ramp from 20% to 85% B between 1 and 25 minutes; a second isocratic hold at 85% B from 25 to 26 minutes; followed by a return to 20% B from 26 to 32 minutes. Filtration of all solvents and solutions was performed using 0.2 μm polyamide membranes (Sartorius Stedim, Goettingen, Germany). An injection of 2 μL was used for the sample. Chromatographic separation was carried out with the column set to 30 $^{\circ}\text{C}$. The electrospray ionization (ESI) source was configured under the following conditions: drying gas heated to 350 $^{\circ}\text{C}$, flow rate of 12 L/min, nebulizer pressure of 55 psi, and a capillary voltage of 4000 V. Additionally, the detection was conducted in dynamic multiple reaction monitoring (dynamic-MRM) mode. The specified peak areas obtained

from dynamic-MRM transitions were used for quantification purposes. The most intense product ion was used for quantitation, while additional fragment ions were used for qualification. A retention time window (Δt) of 2 minutes was applied for each target compound.

2.4 Cell viability assay in tumoral and non-tumoral cells

Three different cell lines were used to analyse cell viability. Human colorectal adenocarcinoma cell line (Caco-2) and human hepatocellular carcinoma cell line (HepG2) were obtained from American Type Culture Collection (ATCC) and cultured in Dulbecco's modified eagle medium (DMEM; Sigma-Aldrich, Barcelona, Spain) enriched with 10% fetal bovine serum (FBS), complemented by 1% penicillin-streptomycin (both from Sigma-Aldrich, Barcelona, Spain). As non-tumoral cell line, the human foreskin fibroblast cell line (HFF-1) from ATCC was selected and cultured with DEMEM, 1% penicillin-streptomycin, 1% pyruvate and 15% FBS. All cell cultures were incubated at 37 °C under a 5% CO₂ atmosphere. Cell viability was determined using thiazolyl blue tetrazolium bromide (MTT) (Fisher Scientific, Barcelona, Spain) by seeding cells in 96-well plates at a concentration of 2×10^4 cells per well and culturing them over a 24-hour period. The accurate quantification of live cells was performed using an automatic cell counter (Logos Biosystems, South Korea) in combination with trypan blue (Sigma-Aldrich, Barcelona, Spain) exclusion staining. Following that, selected wells were exposed to thinned apple by-products within the range of 15.62 to 500 $\mu\text{g mL}^{-1}$, based on previous *in vitro* cytotoxicity studies with thinned apple extracts using similar ranges (Ferrario et al., 2022) (Zheng et al., 2020), prepared in DMEM containing 1% FBS. Control wells received only the medium with 1% FBS, without any extract treatment. After 24 hours, 100 μL of MTT reagent, prepared at 0.4 mg mL^{-1} in a 10% FBS medium, was applied to each well. The plates were then maintained in the dark at 37 °C and 5%

CO₂ environment for 3 hours. After incubation, the MTT solution was removed, and 100 µL of DMSO was introduced into each well to solubilize the formazan precipitate. The resulting purple coloration was indicative of cell viability. The measurement of absorbance was carried out at 550 nm using a Synergy H1 Hybrid Multi-Mode Reader (Biotek, Bad Friedrichshall, Germany). The percentage of viable cells was determined according to the equation below:

$$\text{Cell viability (\%)} = (Abs_{\text{sample}}/Abs_{\text{control}}) \times 100 \quad (1)$$

2.5 *In vitro* antidiabetic assays

2.5.1 α -Glucosidase (from *Saccharomyces cerevisiae*) inhibition

The inhibitory activity of α -glucosidase (Sigma-Aldrich, Barcelona, Spain) was evaluated with a 96-well microplate reader. The α -glucosidase derived from *Saccharomyces cerevisiae* was used as the enzyme source. Each well received 100 µL of α -glucosidase solution (1.0 U mL⁻¹) and 50 µL of the diluted samples at room temperature for a precise preincubation of 10 minutes. Following this, the enzymatic reaction was initiated through the addition of 50 µL of a 3 mM pNPG solution (4-nitrophenyl α -D-glucopyranoside) (Sigma-Aldrich, Barcelona, Spain) as a synthetic substrate. Following a 5-minute incubation at 37 °C to enhance the reaction, the absorbance was recorded at 405 nm. All reagents and extracts were prepared using phosphate buffer. To ensure accurate absorbance readings, control and blank wells were included to eliminate background absorbance and to correct any signal interference caused by the extracts. Acarbose was used as positive control substance. The equation below was applied to quantify the percentage of inhibition:

$$\text{Inhibition (\%)} = [(Abs_{\text{control}} - Abs_{\text{sample}})/Abs_{\text{control}}] \times 100 \quad (2)$$

2.5.2 α -Amylase (from porcine pancreas) inhibition

The inhibition assay of porcine pancreatic α -amylase (Sigma-Aldrich, Barcelona, Spain) was conducted in 1.5 mL microtubes. A volume of 100 μ L of the diluted extract was mixed with 100 μ L of centrifuged α -amylase (2 mg mL⁻¹), followed by a 5-minute incubation at 37 °C. Subsequently, 1% starch solution (Sigma-Aldrich, Barcelona, Spain) was incorporated, and the mixture was kept at 37 °C for an additional 10 min. After that period, 200 μ L of a 1% solution of 3,5-dinitrosalicylic acid (DNS) color reagent (Sigma-Aldrich, Barcelona, Spain) were mixed with 50 μ L of 1 M NaOH, and the mixture was heated at 100 °C for five minutes. Once the reaction concluded, a colour change was observed. Phosphate buffer served as the solvent for all reagents and extract samples. Each reaction mixture (200 μ L), including the control and blanks, was transferred to a 96-well microplate for spectrophotometric analysis at 540 nm. Acarbose served as the positive control substance in the assay. To evaluate inhibitory activity, the following formula was used (2).

2.5.3 Dipeptidyl peptidase-4 (human recombinant DPP-4) inhibition

The commercial DPP-4 inhibitor assay kit (Cayman) was employed to assess the antidiabetic activity of the extracts on the proteolytic activity of human recombinant DPP-4. The assay used the fluorogenic substrate Gly-Pro-Aminomethylcoumarin (AMC), and an incubation of 30 minutes at 37 °C was required to complete the reaction based on the procedure provided by the manufacturer. Fluorescence intensity was monitored employing wavelengths of 350–360 nm for excitation and 450–465 nm for emission. Sitagliptin served as the reference inhibitor. The degree of inhibition was quantified using the equation below.

$$\% \text{ Inhibition} = \left[\frac{(\text{Initial Activity} - \text{Inhibitor})}{\text{Initial Activity}} \right] \times 100 \quad (3)$$

2.6 *In vitro* antiglycation and antioxidant assays

2.6.1 Advanced Glycation End Products (AGEs) inhibition

The accumulation of AGEs was assessed employing a 96-well black microplate format. Each well received 50 μL of sample extract, 100 μL of bovine serum albumin (10 mg mL^{-1}) and 100 μL of fructose (0.5 M), both acquired from Sigma-Aldrich (Barcelona, Spain). All solutions prepared in phosphate buffer. The plate was kept at 37 °C for 24 h in the dark. Control and blanks were included for accurate measurement. Fluorescence intensity was monitored employing wavelengths of 355 nm for excitation and 460 nm for emission. Aminoguanidine was included as the reference inhibitor. To quantify the inhibitory effect, it was represented as a percentage and calculated using equation (2).

2.6.2 Radical superoxide ($\text{O}_2^{\bullet -}$) scavenging from xanthine oxidase system, and xanthine oxidase enzyme inhibition

The neutralization of radical superoxide produced by xanthine oxidase from bovine milk (Sigma-Aldrich, Barcelona, Spain) was evaluated using a 96-well microplate reader. Firstly, a mix solution was prepared with xanthine (Sigma-Aldrich, Barcelona, Spain) (90 μM), nitro tetrazolium blue chloride (NBT) (Sigma-Aldrich, Barcelona, Spain) (22.8 μM) and Na_2CO_3 (Fisher Scientific, Barcelona, Spain) (16 mM) in phosphate buffer. Subsequently, 240 μL of the mixed solution was transferred into each well, followed by 30 μL of the extract in the same buffer and 30 μL of xanthine oxidase (0.168 U mL^{-1}). Control and blanks wells were also prepared for an accurate measurement. After incubation for 2 minutes at 37 °C, absorbance was recorded at 560 nm to determine the radical superoxide scavenging activity of the samples. Quercetin was used as the positive control substance. The percentage of inhibitory activity was calculated according to equation (2).

The inhibition of xanthine oxidase from bovine milk was evaluated in 96-well microplate reader. In this enzymatic inhibition assay, a mix solution was prepared with xanthine (90 μM) and Na_2CO_3 (16 mM) in phosphate buffer. Subsequently, 240 μL of the mixed

solution was added to each well, followed by 30 μL of the extract in the same buffer and 30 μL of xanthine oxidase (0.168 U mL^{-1}). Control and blanks wells were also included. After a 2-minute incubation at 37°C , absorbance was recorded at 295 nm to assess enzymatic inhibition capacity. Inhibition was quantified as a percentage according to equation (2).

2.7 *In vitro* hypolipidemic and hypocholesterolemic assays

2.7.1 Lipase (from porcine pancreas) inhibition

The inhibition of pancreatic lipase (Sigma-Aldrich, Barcelona, Spain) was evaluated through a 96-well plate, with absorbance measured with a microplate reader. Each well contained a reaction mixture composed of 40 μL of diluted extract and 40 μL of centrifuged porcine pancreatic lipase (type II, 2 mg mL^{-1}). The reaction solution was preincubated at room temperature for 15 minutes. Following this, 20 μL of 10 mM p-nitrophenyl butyrate (NPB) (Sigma-Aldrich, Barcelona, Spain) as synthetic substrate was transferred to initiate the reaction, and subsequently the absorbance was recorded at 405 nm and 37°C . Control and blanks wells were also included. All reagents and extracts were prepared using phosphate buffer. Orlistat was utilized as a reference substance for positive control. Lipase inhibition was expressed as a percentage based on the equation (2)

2.7.2 HMG-CoA reductase (purified catalytic domain from HMGR) inhibition

The HMGR inhibitor assay commercial kit (Sigma-Aldrich) was employed to examine the inhibitory activity of the extracts on the HMGR from the purified catalytic domain of the enzyme. The assay relies on a decrease in absorbance at 340 nm caused by the tested extracts and monitors the oxidation of NADPH catalysed by the HMGR enzyme in the presence of its substrate, HMG-CoA. Control and blanks were also used. Based on the manufacturer's instructions, the reaction mixture was held at 37°C for 10 minutes, and

absorbance was recorded at 340 nm. Pravastatin served as the positive control substance. Inhibitory activity was calculated using equation (2).

2.8 *In vitro* hypotensive assays

2.8.1 Angiotensin converting enzyme (ACE from rabbit lung) inhibition

The angiotensin-converting enzyme (ACE) inhibition was evaluated using the ACE commercial kit (Dojindo). This colorimetric bioassay is founded on the enzymatic conversion of the synthetic substrate 3-hydroxybutyryl-Gly-Gly-Gly (3HB-GGG) into 3-hydroxybutyrate (3HB) by ACE. Control and blanks were also performed for an accurate measurement. Following the manufacturer's guidelines, the resulting 3HB was subsequently detected via a coupled reaction that produces a colorimetric signal measurable at 450 nm. Captopril was employed as the positive control substance. Inhibitory activity was calculated using equation (2).

2.9 Statistical Analysis

Statistical procedures were carried out with GraphPad Prism version 10.0 (GraphPad Software, San Diego, CA, USA), including nonlinear regression, one-way ANOVA subsequently analysed using Dunnett's post hoc test, and Pearson correlation analysis. For multivariate data analyse MetaboAnalyst 5.0 (Pang et al., 2021) was carry out as exploratory tool, which included hierarchical clustering heatmaps, principal component analysis (PCA) to enhance comprehension of phenolic content variations. To complement the exploratory PCA visualization, we used PERMANOVA on the corresponding distance matrix to formally test whether overall phenolic profiles differed between cultivars. Statistical significance was assessed using a confidence level of 95% ($p < 0.05$).

Assays were performed in triplicate across separate days, with data reported as mean \pm standard error of the mean (SEM).

2.10 Chemical structure generation

Structures of the main phenolic compounds detected in the extracts were drawn with ChemDraw v21.0.0 (PerkinElmer Informatics, Waltham, MA, USA).

Journal Pre-proof

3. Results

3.1. Phenolic content analyses by HPLC-MS/MS and chemometric analysis

3.1.1. HPLC-MS/MS analysis

In this research, HPLC-MS analysis was performed to target 38 relevant individual phenolic compounds in the 10 thinned apple by-products samples (Table 1). This analytical method identified and quantified 21 distinct phenolic compounds, which included 1 hydroxybenzoic acid, 5 cinnamic acids, 2 dihydrochalcones, 6 flavonols, 4 flavan-3-ols and 3 anthocyanidins. Among them, the greatest diversity of phenolic compounds was identified in the commercial samples of Verde Doncella and Fuji, with 19 compounds detected, while the lowest number, 13 compounds, was found in the autochthonous sample Vadiello-02. Nevertheless, none of the stilbenes nor flavanone compounds were detected in any of the thinned apples.

Furthermore, the highest TPC quantified was detected in the Fuji sample (14010.93 mg kg⁻¹ of extract) followed by Royal Gala, Verde Doncella and Amarilla de Octubre, whereas the lowest TPC was found in the Vadiello-02 sample, with a concentration of 8105.21 mg kg⁻¹ of extract.

The most abundant phenolic compounds quantified in the thinned apple by-products were chlorogenic acid, quercitrin, rutin, hyperoside, isoquercitrin, catechin, procyanidin B2, epicatechin and delphinidin-3,5-diglucoside (Fig.1). Among them, hyperoside, isoquercitrin and delphinidin-3,5-diglucoside were the predominant phenolic compounds in all the samples. In particular, the autochthonous sample Amarilla de Octubre showed the highest concentrations of several compounds, including chlorogenic acid (1945.66 mg kg⁻¹), quercitrin (1567.98 mg kg⁻¹), catechin (502.54 mg kg⁻¹), and procyanidin B2 (1520.56 mg kg⁻¹). Additionally, high levels were recorded for epicatechin in Fuji (1412.45 mg kg⁻¹), rutin in Vadiello-02 (1142.78 mg kg⁻¹), hyperoside in Verde

Doncella (4389.09 mg kg⁻¹), isoquercitrin in Golden Smoothee (4770.77 mg kg⁻¹), and delphinidin 3,5-diglucoside in Royal Gala (4048.11 mg kg⁻¹).

3.1.2. Chemometric analysis

Hierarchical clustering heatmap was used to visualize relative enrichment or depletion of phenolic compounds, and PCA was performed to explore compositional trends between autochthonous and commercial cultivars.

Based on their phenolic profiles, the 10 thinned apple samples were grouped into four distinct clusters (Fig.2). Cluster I grouped the autochthonous samples Daroca-02, Camuesa de Castellano de Daroca, and Vadiello-02, containing, in comparison with the rest of the samples, a higher proportion of ferulic acid, catechin and epicatechin. Cluster II included autochthonous and commercial samples such as Pomera de Pomes Agrias, Reineta, and Golden Smoothee, showing a higher enrichment of phlorizin, neochlorogenic acid, phloretin, and isoquercitrin. Cluster III was formed by the commercial samples Fuji and Royal Gala, which stood out for their higher proportion of quercitrin, hyperoside, p-coumaric acid, phloretin, delphinidin 3- galactoside, delphinidin 3,5-diglucoside, kaempherol-3- glucoside, quercetin, epicatechin, and rutin. Finally, cluster IV consisted of a mixed cultivar sample, such as Amarilla de Octubre and Verde Doncella with a higher proportion of phlorizin, procyanidin B2, gallic acid, catechin, hyperoside, p-coumaric acid, ellagic acid, phloretin, and rutin. Among them, clusters III and IV are notable for exhibiting the highest diversity of phenolic compounds.

The PCA of phenolic groups showed that PC1 and PC2 explained the 46.0% and a 28.9% of total variance, respectively, while the PCA of individual phenolic compounds revealed that PC1 and PC2 explained 28.9% and 25.8% of total variance, respectively. While full class separation is not expected in complex biological matrices the PCA biplot captures most of the phenolic-group variability (74.9% cumulative) and indicates moderate but

statistically significant compositional trends in phenolic-group content across both cultivars ($R^2X = 0.22$; $p = 0.04$). as illustrated in Fig.3. Samples from commercial cultivars are oriented in the direction of the anthocyanins, flavonols, total phenolic content, and total flavonoid content vectors, implying relatively higher concentrations. In contrast, the autochthonous samples are more closely associated with flavan-3-ols, cinnamic acids, dihydrochalcones, and hydroxybenzoic acids, indicating a different phenolic profile. Nevertheless, the PCA analysis of individual phenolic compounds did not reveal statistically significant differences between both cultivars ($R^2X = 0.01$, $p > 0.05$) consistent with the low variance explained by the model.

3.2. Cell viability in Caco-2, HepG2 and HFF-1 cell cultures

The cytotoxic potential of the thinned apple extracts was assessed by exposing two human tumoral cell lines (Caco-2, HepG2) added to a non-tumorigenic human cell line (HFF-1) to a wide concentration range of extracts, from physiologically relevant to non-physiological levels. Following an intestinal–systemic alignment strategy, we used Caco-2 viability to define non-toxic working concentrations and to support intestinal targets, while Caco-2 and HepG2 assays were used to contextualize safe conditions relevant to systemic action (Sanz-Buenhombre et al., 2016) as well as HFF-1 (Liberio et al., 2024). Cell viability was quantified as a percentage, as shown in Fig. 3, with lower viability values indicating higher cytotoxicity. Therefore, to define non-cytotoxic *in vitro* working conditions, the commonly used MTT benchmark (viability $\geq 70\%$) was applied upon exposure of cells to the extracts (Casado-Hidalgo et al., 2025).

Overall, the three cell lines exhibited comparable cytotoxic profiles; however, the tumorigenic HepG2 and Caco-2 cells demonstrated greater sensitivity, particularly at higher concentrations in specific extracts, as evidenced by a more pronounced reduction in viability. The viability reduction induced by some thinned apple extracts followed a

dose dependent pattern across most cell lines, with autochthonous cultivars (Amarilla de Octubre, Pomera de Pomes Agrias, Camuesa de Castellano de Daroca, and Daroca 02) and the commercial extracts (Verde Doncella and Reineta) showing the most pronounced effects at high concentrations.

In HepG2 cells, most of the extracts maintained cell viability above 70% at 500 $\mu\text{g mL}^{-1}$, except for Amarilla de Octubre, Pomera de Pomes Agrias, Reineta, and Verde Doncella. When the concentration was reduced to 250 $\mu\text{g mL}^{-1}$, only Pomera de Pomes Agrias and Amarilla de Octubre remained below this 70% threshold, while at 125 $\mu\text{g mL}^{-1}$, only Amarilla de Octubre showed significant growth inhibition.

A similar trend was observed in Caco-2 cells, where high concentrations of autochthonous (Amarilla de Octubre, Daroca-02, Camuesa Castellano de Daroca) extracts increased cytotoxicity. Specifically, at 500 $\mu\text{g mL}^{-1}$, viability dropped below 70% for the aforementioned cultivars, with Amarilla de Octubre failing to reach this threshold even within the 125–500 $\mu\text{g mL}^{-1}$ range.

In summary, based on viability above 70%, the extracts of Golden Smothee, Fuji, Royal Gala, and Vadiello-02 are classified as non-cytotoxic up to 500 $\mu\text{g mL}^{-1}$ in all cell types assayed. In contrast, Reineta and Verde Doncella are considered safe within physiological ranges ($\leq 250 \mu\text{g mL}^{-1}$ in HepG2, $\leq 500 \mu\text{g mL}^{-1}$ in Caco-2), while Pomera de Pomes Agrias requires concentrations $\leq 125 \mu\text{g mL}^{-1}$ in HepG2 and $\leq 500 \mu\text{g mL}^{-1}$ in Caco-2. Nevertheless Daroca-02 and Camuesa de Castellano de Daroca are considered non-cytotoxic at $\leq 500 \mu\text{g mL}^{-1}$ in HepG2 and $\leq 250 \mu\text{g mL}^{-1}$ in Caco-2. At last, Amarilla de Octubre requires the lowest safe concentrations of all extracts (62.5 $\mu\text{g mL}^{-1}$ in HepG2 and Caco-2) to ensure cytocompatibility.

Conversely, HFF-1 non-tumoral fibroblasts demonstrated higher resilience; even at 500 $\mu\text{g mL}^{-1}$, viability remained between 70–90% for all extracts. This suggests a

selective response, where metabolic function is preserved in non-tumoral human cells. Since the MTT assay measures mitochondrial dehydrogenase activity rather than direct cell lysis, these results likely reflect a metabolic slowdown at supra-physiological concentrations rather than irreversible apoptosis in the analyzed cell types, which is supported by the maintenance of normal cell morphology and adherence observed under microscopic evaluation.

3.3. Evaluation of the *in vitro* antidiabetic activity

3.3.1. Inhibition of α -glucosidase enzyme

A dose-dependent inhibition of α -glucosidase was observed for all extracts. Notably, Amarilla de Octubre ($IC_{50} = 105.60 \pm 7.23 \mu\text{g mL}^{-1}$) exhibited the highest enzymatic inhibition followed by Pomera de Pomes Agrias ($IC_{50} = 123.71 \pm 11.70 \mu\text{g mL}^{-1}$), which showed a lower half-maximal inhibitory concentration (IC_{50}) than the therapeutic agent acarbose ($IC_{50} = 220.00 \pm 64.10 \mu\text{g mL}^{-1}$), indicating greater antidiabetic activity, although without statistically significant difference compared to the control with p values of 0.58 and 0.76, respectively. Furthermore, Vadiello-02 ($p = 0.99$), and Verde Doncella ($p = 0.22$), produced results comparable to the reference drug as well, with IC_{50} values of $244.14 \pm 15.08 \mu\text{g mL}^{-1}$ and $361.56 \pm 6.04 \mu\text{g mL}^{-1}$, respectively, showing no statistically significant differences compared to acarbose ($p > 0.05$). The Reineta extract ($IC_{50} = 486.47 \pm 8.85 \mu\text{g mL}^{-1}$) also stood out above the rest of the samples, showing less significant differences respect to the control ($p = 0.005$), in contrast to the other samples ($p < 0.0001$).

3.3.2. Inhibition of α -amylase enzyme

Only five extracts were capable of inhibiting α -amylase enzyme in a dose-dependent manner. Among them, Vadiello-02 extract exhibited the highest inhibitory activity, followed by Pomera de Pomes Agrias, Amarilla de Octubre, Verde Doncella and Reineta

extracts. Notably, Vadiello-02 ($p = 0.07$) showed an IC_{50} value of $194.79 \pm 22.88 \mu\text{g mL}^{-1}$, which was comparable to the reference antidiabetic drug acarbose ($IC_{50} = 7.72 \pm 1.89 \mu\text{g mL}^{-1}$), with no statistically significant difference observed ($p > 0.05$) compared to control. In addition, Pomera de Pomes Agrias ($IC_{50} = 337.12 \pm 59.73 \mu\text{g mL}^{-1}$) showed great enzymatic inhibition as well, with a less significant difference ($p = 0.002$) compared to control, in contrast to the other samples ($p < 0.0001$).

3.3.3. Inhibition of DPP-4 enzyme

All extracts demonstrated DPP-4 inhibition in a dose-dependent manner (Fig.4A), contributing to the overall synergistic antidiabetic potential of the samples, although with generally low inhibitory potency against DPP-4. Among these, Pomera de Pomes Agrias ($IC_{50} = 480.63 \pm 3.39 \mu\text{g mL}^{-1}$), followed by Amarilla de Octubre ($IC_{50} = 628.03 \pm 21.09 \mu\text{g mL}^{-1}$), demonstrated the highest inhibitory activity. Both showed less pronounced statistically significant differences compared to the reference antidiabetic drug sitagliptin ($IC_{50} = 0.04 \pm 0.01 \mu\text{g mL}^{-1}$), with p -values of 0.003 and 0.0001, respectively, in contrast to the other samples ($p < 0.0001$). These results emphasize the antidiabetic potential of Pomera de Pomes Agrias and Amarilla de Octubre extracts.

3.4. Evaluation of the *in vitro* assay of the antiglycant and antioxidant activity

3.4.1. Inhibition of AGEs formation

In this non-enzymatic assay, all extracts demonstrated the ability to inhibit albumin protein glycation. Specifically, Amarilla de Octubre and Pomera de Pomes Agrias extracts, with IC_{50} values of $49.77 \pm 3.52 \mu\text{g mL}^{-1}$ and $60.08 \pm 3.01 \mu\text{g mL}^{-1}$, respectively, exhibited the most pronounced antiglycation effects. The Reineta sample ($IC_{50} = 189.36 \pm 13.15 \mu\text{g mL}^{-1}$), followed by Verde Doncella ($IC_{50} = 203.44 \pm 5.61 \mu\text{g mL}^{-1}$) also stood out in terms of antiglycation bioactivity. It is noteworthy that Amarilla de Octubre ($p = 0.99$), exhibited stronger activity, as indicated by a lower IC_{50} value, whereas Pomera de

Pomes Agrias ($p = 0.99$) showed a higher IC_{50} than the investigational drug aminoguanidine ($IC_{50} = 55.01 \pm 5.63 \mu\text{g mL}^{-1}$). Both extracts showed notable antiglycation effects without statistically significant differences compared to the control ($p > 0.05$), in contrast to the remaining samples ($p < 0.0001$).

3.4.2. Antioxidant activity against radical superoxide ($O_2^{\bullet -}$) and xanthine oxidase

This assay allowed us to differentiate the antioxidant mechanisms of action between superoxide radical scavenging and xanthine oxidase enzyme inhibition. A clear antioxidant mechanism was identified, as all extracts demonstrated superoxide radical scavenging activity, while none of the extracts were able to inhibit the enzyme. The antioxidant reference substance was the pure phenolic compound quercetin ($IC_{50} = 4.16 \pm 0.24 \mu\text{g mL}^{-1}$), which stood out due to its strong radical scavenging ability. Nevertheless, the extracts of Amarilla de Octubre ($IC_{50} = 1.18 \pm 0.21 \mu\text{g mL}^{-1}$) followed by Pomera de Pomes Agrias ($IC_{50} = 1.98 \pm 0.02 \mu\text{g mL}^{-1}$), Fuji ($IC_{50} = 2.13 \pm 0.29 \mu\text{g mL}^{-1}$), Reineta ($IC_{50} = 2.47 \pm 0.50 \mu\text{g mL}^{-1}$) and Verde Doncella ($IC_{50} = 2.75 \pm 0.26 \mu\text{g mL}^{-1}$) achieved even higher scavenging activity than the control substance. All extracts showed comparable antioxidant activity to quercetin, with no statistically significant differences observed ($p > 0.05$), except for the Vadiello-02 extract ($p < 0.001$), which exhibited the lowest antioxidant capacity.

3.5. Evaluation of the *in vitro* hypolipidemic and hypocholesterolemic activity

3.5.1. Inhibition of lipase enzyme

All extracts showed dose-dependent inhibition of lipase activity. Notably, the Reineta extract ($IC_{50} = 225.43 \pm 6.31 \mu\text{g mL}^{-1}$), followed by the Verde Doncella extract ($IC_{50} = 363.90 \pm 25.91 \mu\text{g mL}^{-1}$), Fuji ($IC_{50} = 397.1 \pm 44.77 \mu\text{g mL}^{-1}$), Amarilla de Octubre ($IC_{50} = 419.31 \pm 17.76 \mu\text{g mL}^{-1}$) and Pomera de Pomes Agrias ($IC_{50} = 447.02 \pm 25.45 \mu\text{g mL}^{-1}$) showed the most pronounced hypolipidemic effects. These extracts exhibited statistically

significant differences compared to the reference hypolipidemic agent orlistat ($IC_{50} = 27.12 \pm 11.49 \mu\text{g mL}^{-1}$). However, the Reineta extract ($p = 0.03$) and the Verde Doncella extract ($p = 0.0001$) showed less pronounced statistical differences than the other samples ($p < 0.0001$), indicating a closer similarity to the control treatment.

3.5.2. Inhibition of HMGR enzyme

In this enzymatic assay, only four extracts were able to inhibit HMGR activity (Fig. 4B), acting through a more specific and complex mechanism of action in HMGR enzyme in a dose-dependent manner. Among them, Verde Doncella ($IC_{50} = 100.88 \pm 3.93 \mu\text{g mL}^{-1}$) exhibited the highest hypocholesterolemic activity, followed by Reineta ($IC_{50} = 149.39 \pm 9.39 \mu\text{g mL}^{-1}$), Vadiello-02 ($IC_{50} = 233.04 \pm 17.78 \mu\text{g mL}^{-1}$) and Golden Smoothee ($IC_{50} = 469.49 \pm 8.04 \mu\text{g mL}^{-1}$) extracts. In particular, Verde Doncella extract exhibited less pronounced statistically significant differences ($p = 0.0001$) compared to the reference hypocholesterolemic agent pravastatin ($IC_{50} = 0.012 \pm 0.0001 \mu\text{g mL}^{-1}$), in contrast to the other thinned apple extracts ($p < 0.0001$). These findings emphasize the potential of Verde Doncella and Reineta extracts as promising hypolipidemic and hypocholesterolemic agents, acting through a synergistic mechanism of action.

3.6. Evaluation of the *in vitro* activity against ACE enzyme

All extracts were able to inhibit the ACE enzyme (Fig. 4C), showing an additional hypotensive activity. Among them, Verde Doncella ($IC_{50} = 42.17 \pm 3.11 \mu\text{g mL}^{-1}$) followed by Amarilla de Octubre ($IC_{50} = 55.94 \pm 1.18 \mu\text{g mL}^{-1}$), Pomera de Pomes Agrias ($IC_{50} = 81.32 \pm 10.33 \mu\text{g mL}^{-1}$), Reineta ($IC_{50} = 84.34 \pm 11.58 \mu\text{g mL}^{-1}$) and Daroca-02 ($IC_{50} = 90.91 \pm 12.09 \mu\text{g mL}^{-1}$) showed the highest ACE inhibition. Furthermore, Verde Doncella ($p = 0.26$) and Amarilla de Octubre ($p = 0.07$) showed no statistically significant differences ($p > 0.05$) compared to the hypotensive agent captopril ($IC_{50} = 0.003 \pm 0.001 \mu\text{g mL}^{-1}$), revealing a similarly interesting hypotensive activity. In addition, the samples

of Pomera de Pomes Agrias ($p = 0.004$), Reineta ($p = 0.003$), and Daroca-02 ($p = 0.001$), exhibited a notable capacity to inhibit the ACE enzyme, showing less statistically significant differences ($p < 0.05$) compared to captopril, in contrast to the remaining extracts which displayed more pronounced differences ($p < 0.0001$).

3.7. Determination of correlation between all *in vitro* bioassays and bioassays with individual phenolic content.

The analysis of the IC_{50} of the enzymes revealed several statistically significant positive correlations, suggesting potential functional relationships among metabolic and regulatory pathways. The strongest positive correlation was observed between AGEs and radical superoxide scavenging activity ($R = 0.92$, $p = 0.0001$), indicating a close link between oxidative stress and glycation processes. Other notable positive correlations include ACE with AGEs ($R = 0.82$, $p = 0.004$), as well as ACE with radical superoxide scavenging ($R = 0.71$, $p = 0.02$). These findings highlight possible interactions between oxidative stress, glycation, and blood pressure regulation mechanisms. Similarly, α -glucosidase and α -amylase exhibited a strong positive association ($R = 0.88$, $p = 0.0007$), consistent with their complementary roles in carbohydrate digestion. A moderate correlation was also found between α -glucosidase and DPP-4 ($R = 0.65$, $p = 0.04$), reinforcing the interplay in glucose metabolism.

In addition, several phenolic compounds demonstrated strong and statistically significant negative correlations with the IC_{50} of the enzymes, as shown in Fig. 5, suggesting potential inhibitory effects and relevant bioactivity. Significant negative correlations were identified for AGEs with procyanidin B2 ($R = -0.72$, $p = 0.02$), gallic acid ($R = -0.70$, $p = 0.02$), and phlorizin ($R = -0.65$, $p = 0.04$), indicating potential roles in reducing advanced glycation end-products. Furthermore, radical superoxide scavenging exhibited negative associations with phlorizin ($R = -0.69$, $p = 0.02$), procyanidin B2 ($R = -0.64$, p

= 0.04), and phloretin ($R = -0.64$, $p = 0.04$), while gallic acid ($R = -0.78$, $p = 0.008$) together with procyanidin B2 ($R = -0.86$, $p = 0.001$) with ACE, reinforcing their cardiovascular relevance, with procyanidin B2 being the most pronounced inverse association. These mechanisms exhibited multiple significant negative correlations with different phenolics, suggesting synergistic inhibitory effects and emphasizing the importance of polyphenol synergy and features in enhancing enzyme inhibition and improving cardiometabolic health.

Additionally, quercitrin showed a strong negative correlation with DPP-4 ($R = -0.83$, $p = 0.003$), highlighting its possible contribution to glycemic control. Other notable findings include phloretin with lipase ($R = -0.79$, $p = 0.006$), suggesting an effect on lipid digestion, also HMGR correlated negatively with isoquercitrin ($R = -0.65$, $p = 0.043$), suggesting possible implications for cholesterol metabolism. Nevertheless, α -glucosidase ($R = -0.59$) and α -amylase ($R = -0.55$) enzymes showed moderate negative correlations with gallic acid, but none reached statistical significance ($p > 0.05$) indicating a potential mechanistic trend toward inhibition of carbohydrate digestion without strong evidence under current conditions.

Furthermore, there was no significant correlation ($p > 0.05$), between any of the enzymes and TPC nor TFC of the targeted phenolic compounds. However, TPC appears to be biologically relevant for the antioxidant activity of the extracts in the scavenging of superoxide radicals, as indicated by the observed correlation ($R = -0.51$). However, given that only a partial phenolic analysis was performed, the absence of significant differences is understandable.

Collectively, these findings support the hypothesis that specific polyphenols, particularly procyanidin B2, quercitrin, gallic acid, phloretin, phlorizin, and isoquercitrin may exert beneficial effects by inhibiting enzymes involved in hypertension, diabetes, oxidative

stress and lipid metabolism. Nevertheless, other phenolic compounds showed some degree of correlation with enzymatic inhibition, although without statistical significance, as this could bias enzymes associated with a single phenolic compound. However, their potential contribution to the overall inhibitory biological effect cannot be ruled out, since enzyme inhibition by complex plant matrices often results from synergistic or additive interactions among multiple bioactive compounds. Due to the limited sample size along with the limited bioactive compounds analysed, it is not possible to determine the actual impact of these secondary correlations.

4. Discussion

By delving deeper into the targeted analysis of the phenolic profile using HPLC/MS along with the chemometric analysis, the highest TPC was found in the commercial cluster of Fuji and Royal Gala, followed by the cluster of Verde Doncella and Amarilla de Octubre. This result contrasts with other previous studies focused on the phenolic content of pulp (Millán-Laleona et al., 2025) and peel (Cano-Lou et al., 2025) from different apple cultivars, where autochthonous samples were richer in the TPC than the commercial samples. This could be explained by the rise of sugar content, added to a reduction in phenolic compounds and the antioxidant activity during the ripening process, especially in commercial samples (Zheng et al., 2012).

As shown, all thinned apple by-products are rich sources of phenolic compounds such as hyperoside, isoquercitrin, and delphinidin-3,5-diglucoside, followed by chlorogenic acid, quercitrin, rutin, catechin, procyanidin B2, and epicatechin with yields ranging between 20-30% (w/w) of dry ethanolic extracts relative to the weight of the lyophilized samples and between 8105.21 to 14010.93 mg kg⁻¹ of TPC. Indicating that bulk extract recovery does not necessarily translate into proportional differences in phenolic loading. Additionally, the great yield as well as the rich phenolic matrix obtained suggests that this

type of residue is a great source of phenolic compounds, and that the extraction method along with the ethanol solvent system is efficient (Do et al., 2014).

The PCA biplot analysis suggests compositional trends in phenolic group content, with a higher ratio of anthocyanins and flavonols in commercial samples, and a greater presence of flavan-3-ols, cinnamic acids, dihydrochalcones, and hydroxybenzoic acids in autochthonous samples. These results align with previous research comparing commercial and autochthonous apple peel by-products (Cano-Lou et al., 2025).

Regarding the PCA of individual phenolic compounds and the hierarchical clustering heatmap, both indicate that autochthonous and commercial samples display high variability in phenolic distribution, with no clear separation and mixed clusters containing both commercial and autochthonous cultivars. Differences in phenolic composition among cultivar clusters clearly contributed to the wide range of bioactivities observed across the thinned apple extracts against multiple MetS enzymatic targets. Within all samples, cluster II of the hierarchical clustering heatmap, especially Reineta and Pomera de Pomes Agrias, as well as cluster IV of Verde Doncella-Amarilla de Octubre exhibited the highest results in all the bioactivities performed, pointing out the relationship between their individual phenolic content and the great bioactive properties. However, the autochthonous samples of both clusters (Pomera de Pomes Agrias, Amarilla de Octubre) exhibited higher antioxidant, antiglycant and antidiabetic activity, in contrast to the commercial samples of both clusters (Reineta, Verde Doncella) which exhibited greater hypolipidemic and hypocholesterolemic activity. In addition, cluster IV exhibited the greatest hypotensive activity. Apart of these samples, Vadiello-02 from all autochthonous cluster I also showed great antidiabetic and hypocholesterolemic activity, despite being the sample with the lowest TPC.

These findings suggest that the inhibitory activity seems to be more strongly associated with the specific molecular structures of the phytochemicals and the conformational properties of the target proteins, which govern the formation of protein–phenolic complexes essential for inhibition (Dalar & Konczak, 2013). The evidence presented indicates that the targeted TPC is not a reliable predictor of enzyme inhibitory capacity of thinned apple extracts. Depending on their specific chemical structures such as the quantity and positional distribution of hydroxyl groups, the glycosylation and the structural complexity of phenolic compounds may influence the interaction with the enzymatic catalytic site of digestive enzymes (Martinez-Gonzalez et al., 2017), DPP-4 (Huang et al., 2019), ACE (Chen et al., 2021) and HMGR through hydrophobic interactions, metal chelation, or hydrogen bonding. However, in HMGR, steric limitations often redirect binding to the NADPH cofactor site (Ojha et al., 2015) making it a unique exception among these enzymes, which may explain why only 4 out of 10 tested extracts exhibited inhibitory activity against this enzyme.

Consequently, the proportion of individual phenolic compounds and their interactions may significantly influence the antioxidant activity, as well as the antidiabetic, hypotensive, hypolipidemic, and hypocholesterolemic effects (Yeh et al., 2017) of thinned apple by-products on MetS.

For instance, procyanidin B2 and phlorizin were present at higher levels in clusters II and IV than the other cultivars, and their significant contribution to radical-scavenging capacity and AGEs inhibition, together with elevated gallic acid concentrations as well, is consistent with enhanced antiglycation potential. This compositional interplay may explain why these autochthonous and commercial clusters outperform the remaining samples in these specific activities. Specifically, Amarilla de Octubre showed the strongest antioxidant and antiglycation activities, which coincided with the highest

concentrations of procyanidin B2 (1520.56 mg kg⁻¹), gallic acid (23.11 mg kg⁻¹), and phlorizin (323.67 mg kg⁻¹). The co-enrichment of phlorizin, procyanidin B2, and gallic acid in the most active extracts is consistent with prior studies that report antioxidant and antiglycation activities of these individual phenolic compounds (Chen et al., 2023; Bai et al., 2021; Maisto et al., 2022).

Furthermore, quercitrin was the only phenolic compound significantly associated with an antidiabetic *in vitro* bioactivity, showing a strong significant correlation with DPP-4 inhibition and reaching its highest levels in Amarilla de Octubre and Pomera de Pomes Agrias (1567.98 and 1447.40 mg kg⁻¹). Notably, these two extracts also exhibited the highest DPP-4 inhibitory activity, supporting quercitrin as a key contributor to glycaemic regulation. Likewise, the enrichment in quercitrin can reasonably support DPP-4 inhibition, since isolated quercitrin (quercetin-3-O-rhamnoside) inhibited purified DPP-4 (from Cayman) in a biochemical assay with an IC₅₀ of 71.79 ± 1.05 µg mL⁻¹ (Liu et al., 2021).

In addition, isoquercitrin and phloretin emerged as the phenolic features most aligned with lipid related *in vitro* activities. Isoquercitrin was particularly abundant in Verde Doncella and Reineta (3550.87 and 3755.56 mg kg⁻¹) and showed a significant correlation with HMGR activity, supporting a potential contribution to hypocholesterolemic effects via modulation of cholesterol biosynthesis. In addition, phloretin, detected in clusters II and IV, correlated strongly with the lipase inhibition, and despite its very low abundance it may still contribute to the activity pattern, particularly in Reineta and Verde Doncella. The enrichment of isoquercitrin in the extracts showing stronger HMGR effects are biologically plausible because isolated isoquercitrin (quercetin-3-O-glucoside) has been shown to inhibit purified HMGR (from Sigma Aldrich) in a spectrophotometric assay with an IC₅₀ of 80.6 µM (Kwon et al., 2010) corresponding to 37.4 µg mL⁻¹. In addition,

the enrichment of the extracts in phloretin and their bioactivity in inhibiting pancreatic lipase are confirmed by the inhibition of the enzyme by pure phloretin, which exhibits an IC_{50} of $164.7 \pm 2.4 \mu\text{M}$ (Jeong et al., 2021) corresponding to $45.2 \mu\text{g mL}^{-1}$.

Moreover, procyanidin B2 emerged as the key phenolic driver of ACE inhibition, showing the strongest significant correlation and reaching high levels in Amarilla de Octubre ($1520.56 \text{ mg kg}^{-1}$) and Verde Doncella ($1456.08 \text{ mg kg}^{-1}$), among the highest across all samples. Consistently, these two varieties displayed the most pronounced ACE inhibitory activity, supporting procyanidin B2 as a major contributor to renin–angiotensin system modulation. Procyanidin B2 is known as the most effective ACE inhibitor among plant flavonoids, as epicatechin dimers have been shown to inhibit ACE (from rabbit lung) with an IC_{50} of $97 \mu\text{M}$ (Nileeka Balasuriya & Vasantha Rupasinghe, 2011), corresponding to $56.1 \mu\text{g mL}^{-1}$.

The IC_{50} value of the most bioactive extracts was within the same order of magnitude as that IC_{50} reported for the previous isolated phenolic compounds. Considering that the previous pure compounds are present in the thinned apple extracts at a lower concentration, this comparable activity suggest that weaker active constituents may contribute to the observed bioactivity (Atanasov et al., 2015). Whether these interactions are additive or synergistic cannot be determined from the present data and warrants further investigation through combination studies (Caesar & Cech, 2019). Furthermore, the lack of significant correlation between antidiabetic digestive enzymes and the targeted phenolics indicates that α -glucosidase and α -amylase inhibition may depend on specific phenolic mixtures or non-targeted metabolites. For instance, lipophilic triterpenes (ursolic and oleanolic acids), reported in apple matrices and linked to carbohydrate enzyme inhibition, may contribute alongside phenolic mixtures in the most active extracts

(Amarilla de Octubre, Pomera de Pomes Agrias, Vadiello-02) (Andre et al., 2012; Hussain et al., 2017).

Accordingly, to ensure that the observed bioactivity is interpreted within a cytocompatible window, we evaluated extract-induced cytotoxicity across representative structural, hepatic, and intestinal models as phenolic compounds can induce some disrupting mitochondrial functions (Procházková et al., 2011).

Most of the thinned apple extracts exhibited a robust safety profile at physiological exposure levels ($\geq 70\%$ viability) in non-tumorigenic cells. HFF-1 non-tumorigenic cells were the most resilient model, maintaining the defined cytocompatibility across the full tested range for all extracts ($15.62\text{--}500\ \mu\text{g mL}^{-1}$). By comparison, HepG2 and Caco-2 cells, which are tumorigenic models, showed moderate tolerance for most cultivars, but a reduction in viability was detected particularly in Amarilla de Octubre and Pomera de Pomes Agrias, which are enriched in specific antiproliferative phenolic compounds such as quercitrin (Cincin et al., 2014) and chlorogenic acid (Jiang et al., 2021). This fact could explain the cytotoxic/antiproliferative activity of certain extracts in these hepatic and colon cancer-related cell lines.

Collectively, these cell-type and cultivar-dependent thresholds suggest different upper tolerance boundaries under *in vitro* exposure conditions, with fibroblasts exhibiting the widest safety window and intestinal-hepatic models revealing cultivar-specific sensitivity. These findings agree with previous studies on ethanolic thinned-apple polyphenol extracts, which reported significant antioxidant and anti-inflammatory *in vitro* activities with minimal cytotoxicity at comparable doses, maintaining viability $\geq 70\%$ up to $250\text{--}300\ \mu\text{g mL}^{-1}$ in HEK293 and R3/1-NF- κ B cells (Ferrario et al., 2022), as well as in murine macrophage-like cells (Zheng et al., 2020). This indicates that *in vitro* bioactivity can be achieved within defined non-cytotoxic windows.

The increased sensitivity observed in HepG2 and Caco-2 tumoral cell lines at high concentrations such as 500 $\mu\text{g mL}^{-1}$ for specific cultivars (Amarilla de Octubre, Pomera de Pomes Agrias, Camuesa de Castellano de Daroca, Daroca-02, Reineta and Verde Doncella) is consistent with dose-dependent cytotoxicity reported of thinned apple extracts, where tumor-derived murine macrophage cells showed a viability decline to 50% at 600 $\mu\text{g mL}^{-1}$ (Zheng et al., 2020). This dose-responsive loss of viability supports the notion that, although apple-derived complexes are generally well tolerated, certain cultivar-specific compositions may approach cellular stress thresholds at certain concentrations. For instance, the samples that induced the strongest reduction in HepG2 cell viability at lower concentrations (Amarilla de Octubre and Pomera de Pomes Agrias) were those most enriched in chlorogenic acid, a phenolic compound highly associated with enhanced antiproliferative activity in hepatic tumor models (Jiang et al., 2021).

In addition, HepG2 hypersensitivity should be interpreted cautiously because *in vivo* hepatocyte exposure occurs only after intestinal absorption and first-pass metabolism, resulting in substantially lower exposition concentrations than those applied in *in vitro* assays. In contrast, the generally higher tolerance observed in Caco-2 supports the feasibility of local intestinal applications, where luminal concentrations higher than systemic levels may be required for efficacy without compromising epithelial viability within defined cytocompatible ranges (Sahu et al., 2014).

Moreover, *in vitro* IC_{50} values of thinned apple extracts should be interpreted in the context of achievable exposure, ideally by benchmarking them against unbound plasma concentrations and target tissue levels (Abdel-Tawab, 2021). Considering that phenolic compounds typically circulate at relatively low concentrations (10–500 ng mL^{-1}), direct systemic inhibition may be modest, whereas higher local tissue levels (up to 1 $\mu\text{g mL}^{-1}$) can make target modulation more plausible (Yoshida et al., 2017). In this setting, efficacy

may be further enhanced by the combined action of metabolites, tissue accumulation, and site-specific exposure such as in hepatic parenchyma for HMGR, vascular endothelium for ACE, and multiple systemic tissues for DPP-4 (Abdel-Tawab, 2021). Crucially, these concentrations remain within the safety boundaries of Caco-2, HepG2 and HFF-1 cells enhancing metabolic modulation.

The multitarget effect towards different enzymes (ACE, HMGR), combined with antiglycan activity and a potent antioxidant protection towards $O_2^{\bullet-}$ scavenging, may yield meaningful benefits by acting on several pathways simultaneously. Specifically, Verde Doncella and Reineta demonstrated a superior therapeutic window for cardiovascular and lipid management, inhibiting ACE and HMGR below $250 \mu\text{g mL}^{-1}$ under non-cytotoxic conditions. Additionally, Amarilla de Octubre and Pomera de Pomes Agrias also contributed to ACE, AGEs inhibition and radical scavenging at concentrations that can be considered cytocompatible, particularly in the non-tumoral HFF-1 cells. The most biological relevance of these thinned apple extracts lies in the correlation of antioxidant, anti-glycation, and ACE-inhibitory bioactivities together with an enriched profile of bioactive compounds such as procyanidin B2, gallic acid and phlorizin. This convergence suggests a coherent chemical feature underlying the observed functional effects.

Notably, several extracts demonstrated *in vitro* bioactivities comparable to reference substances with no statistically significant differences ($p > 0.5$) in AGEs inhibition (Pomera de Pomes Agrias, Amarilla de Octubre), ACE inhibition (Verde Doncella, Amarilla de Octubre), and antioxidant activity (all extracts except Vadiello-02); such bioactivities reinforce the potential translational value of these extracts. In contrast, HMGR inhibition was considered of lower biological relevance because it was only detected in four extracts (Reineta, Verde Doncella, Golden Smoothee and Vadiello-02),

associated with a single phenolic marker (isoquercitrin), and markedly less potent than pravastatin. Meanwhile, the DPP-4 inhibition was not considered interesting as IC_{50} values were very high. Within this framework, Clusters II and IV, and particularly Amarilla de Octubre, emerge as key candidates for biological relevance in MetS. This claim is supported by the highest contain in procyanidin B2, gallic acid, and phlorizin, which, despite representing only 0.19% of the total extract mass in Amarilla de Octubre, potentially points toward a high antiglycation and antioxidant activity along with inhibitory interaction with ACE.

In addition, the cultivar Pomera de Pomes Agrias also stand out as a prominent multitarget inhibitor in key intestinal enzymes involved in carbohydrate metabolims (α -glucosidase, α -amylase, DPP-4) while maintaining IC_{50} values under non-cytotoxic conditions in Caco-2. This activity profile supports their biological relevance as candidate for the development of nutraceuticals with antidiabetic properties.

Furthermore, the obtained data encourage deeper investigation for potential industrial applications and incorporation of isolated compounds or complex mixtures into nutraceuticals and dietary supplements in the health and well-being sector.

Conclusion

Targeted HPLC–MS profiling of UAE ethanolic extracts from thinned apples identified hyperoside, isoquercitrin, and delphinidin-3,5-diglucoside as the predominant phenolics across samples. The marked cultivar-driven diversity supported the efficiency of the extraction approach, and PCA clearly separated commercial cultivars from autochthonous cultivars phenolic groups. However, hierarchical clustering at the level of individual phenolics produced mixed clusters containing both origins, indicating clusters enriched in specific phenolic compounds. Integrating these chemical targeted fingerprints with

chemometric modelling linked composition to biological potency (IC_{50}), enabling early prioritization of predictive metabolite features and supporting a data-guided selection of candidate extracts or enriched mixtures for health applications. Accordingly, phenolic compounds such as procyanidin B2, quercitrin, gallic acid, phloretin, phlorizin, and isoquercitrin emerge as plausible contributors to multi-pathway metabolic modulation supported by significant correlation.

In addition, the cultivars Amarilla de Octubre and Pomera de Pomes Agrias showed the best antioxidant-antiglycation activity, with interesting antidiabetic activity (due to α -glucosidase, α -amylase and DPP-4 inhibition) while Reineta and Verde Doncella showed the most pronounced hypolipidemic/cardiovascular activities (due to lipase, HMGR and ACE inhibition). Such findings highlight a possibility to use thinned apples as food matrices for the extraction of phenolic compounds and development of nutraceuticals, emphasizing the value of reusing by-products generated by the agronomy sector for new industrial applications.

Data availability

The authors confirm that all relevant data supporting the findings of this research are included in the manuscript. Additional raw and processed datasets can be obtained from the corresponding author upon reasonable request.

Conflicts of interest

There are no conflicts to declare.

Funding

This work is part of the APPLIEDIV Project (Multidisciplinary approach for exploiting genetic diversity to increase the value of autochthonous apple: valorizing local apple by-products and wastes through bioactive compounds PID2022–141847OR-C33) funded by

the Research State Agency in the 2022 call for Projects of Generation Knowledge oriented towards societal challenges.

Acknowledgements

The authors thank Teva Pharma S.L.U for the research PhD grant to Javier Cano-Lou, the Government of Aragon for financial support from the Phytopharm group (ref. B44_23R) and Universidad San Jorge for funding Internal Project 2425013 through the Call for Internal Research Projects (Academic Year 2024–2025).

Abbreviations

ACE Angiotensin-converting enzyme inhibitors

AGEs Advanced glycation end products

Caco-2 Human colorectal adenocarcinoma cell line

CVD Cardiovascular disease

DMEM Dulbecco's modified eagle medium

DNS Dinitrosalicylic acid

DPP-4 Dipeptidyl peptidase-4

ESI Electrospray ionization

FBS Foetal bovine serum

GAE Gallic acid equivalent

HDL High-density lipoprotein

HepG-2 Cells human hepatocellular carcinoma cell line

HFF-1 Human foreskin fibroblast cell line

HMGR 3-hydroxy-3-methylglutaryl-CoA reductase

HPLC-MS/MS High performance liquid chromatography coupled with mass spectrometry

IC₅₀ Half-maximal inhibitory concentration

MetS Metabolic syndrome

MTT Thiazolyl blue tetrazolium bromide

NPB p-4-Nitrophenyl butyrate

PCA Principal component analysis

pNPG 4-nitrophenyl α -D-glucopyranoside

ROS Reactive oxygen species

SEM Standard error of mean

SGLTs Sodium-dependent glucose transporters

SLR Solid–liquid ratio

T2D Type 2 diabetes

TPC Total phenolic content

UAE Ultrasound-assisted extraction

Journal Pre-proof

5. References

1. Abdel-Tawab, M. (2021). Considerations to Be Taken When Carrying Out Medicinal Plant Research—What We Learn from an Insight into the IC₅₀ Values, Bioavailability and Clinical Efficacy of Exemplary Anti-Inflammatory Herbal Components. *Pharmaceuticals*, *14*(5), 437. <https://doi.org/10.3390/ph14050437>
2. Andre, C. M., Greenwood, J. M., Walker, E. G., Rassam, M., Sullivan, M., Evers, D., Perry, N. B., & Laing, W. A. (2012). Anti-Inflammatory Procyanidins and Triterpenes in 109 Apple Varieties. *Journal of Agricultural and Food Chemistry*, *60*(42), 10546-10554. <https://doi.org/10.1021/jf302809k>
3. Atanasov, A. G., Waltenberger, B., Pferschy-Wenzig, E.-M., Linder, T., Wawrosch, C., Uhrin, P., Temml, V., Wang, L., Schwaiger, S., Heiss, E. H., Rollinger, J. M., Schuster, D., Breuss, J. M., Bochkov, V., Mihovilovic, M. D., Kopp, B., Bauer, R., Dirsch, V. M., & Stuppner, H. (2015). Discovery and resupply of pharmacologically active plant-derived natural products: A review. *Biotechnology Advances*, *33*(8), 1582-1614. <https://doi.org/10.1016/j.biotechadv.2015.08.001>
4. Bai, J., Zhang, Y., Tang, C., Hou, Y., Ai, X., Chen, X., Zhang, Y., Wang, X., & Meng, X. (2021). Gallic acid: Pharmacological activities and molecular mechanisms involved in inflammation-related diseases. *Biomedicine & Pharmacotherapy*, *133*, 110985. <https://doi.org/10.1016/j.biopha.2020.110985>
5. Caesar, L. K., & Cech, N. B. (2019). Synergy and antagonism in natural product extracts: When 1 + 1 does not equal 2. *Natural Product Reports*, *36*(6), 869-888. <https://doi.org/10.1039/C9NP00011A>
6. Cano-Lou, J., Millán-Laleona, A., Candrea, R., Les, F., Pina, A., Caprioli, G., & López, V. (2025). Apple peels as an edible source of phenolic bioactive compounds

- with antidiabetic and antiglycation properties. *Food & Function*, 16(8), 2947-2958.
<https://doi.org/10.1039/d4fo05241b>
7. Capanoglu, E., Nemli, E., & Tomas-Barberan, F. (2022). Novel Approaches in the Valorization of Agricultural Wastes and Their Applications. *Journal of Agricultural and Food Chemistry*, 70(23), 6787-6804. <https://doi.org/10.1021/acs.jafc.1c07104>
 8. Casado-Hidalgo, G., Cebollada, P., Cano-Lou, J., Cardoso, R. V. C., Barros, L., Rodríguez-Yoldi, M. J., & López, V. (2025). Boletus edulis as a healthy and prized edible mushroom: Analysis of bioactive compounds and in vitro functional properties. *Applied Food Research*, 5(2), 101342. <https://doi.org/10.1016/j.afres.2025.101342>
 9. Chen, J., Zhong, K., Jing, Y., Liu, S., Qin, S., Peng, F., Li, D., & Peng, C. (2023). Procyanidin B2: A promising multi-functional food-derived pigment for human diseases. *Food Chemistry*, 420, 136101. <https://doi.org/10.1016/j.foodchem.2023.136101>
 10. Chen, L., Wang, L., Shu, G., & Li, J. (2021). Antihypertensive Potential of Plant Foods: Research Progress and Prospect of Plant-Derived Angiotensin-Converting Enzyme Inhibition Compounds. *Journal of Agricultural and Food Chemistry*, 69(18), 5297-5305. <https://doi.org/10.1021/acs.jafc.1c02117>
 11. Cincin, Z. B., Unlu, M., Kiran, B., Bireller, E. S., Baran, Y., & Cakmakoglu, B. (2014). Molecular Mechanisms of Quercitrin-induced Apoptosis in Non-small Cell Lung Cancer. *Archives of Medical Research*, 45(6), 445-454. <https://doi.org/10.1016/j.arcmed.2014.08.002>
 12. Dalar, A., & Konczak, I. (2013). Phenolic contents, antioxidant capacities and inhibitory activities against key metabolic syndrome relevant enzymes of herbal teas from Eastern Anatolia. *Industrial Crops and Products*, 44, 383-390. <https://doi.org/10.1016/j.indcrop.2012.11.037>

13. Do, Q. D., Angkawijaya, A. E., Tran-Nguyen, P. L., Huynh, L. H., Soetaredjo, F. E., Ismadji, S., & Ju, Y.-H. (2014). Effect of extraction solvent on total phenol content, total flavonoid content, and antioxidant activity of *Limnophila aromatica*. *Journal of Food and Drug Analysis*, 22(3), 296-302. <https://doi.org/10.1016/j.jfda.2013.11.001>
14. Dou, Z., Tian, Y., Zhang, Y., Zhang, W., Duan, Q., Huang, Q., Zhang, B., Niu, H., Wang, L., & Zeng, S. (2025). Pectin-type polysaccharides from raspberry (*Rubus idaeus* L.): Structure characterization and activity against DSS-induced colitis. *Carbohydrate Polymers*, 364, 123710. <https://doi.org/10.1016/j.carbpol.2025.123710>
15. Fahed, G., Aoun, L., Bou Zerdan, M., Allam, S., Bou Zerdan, M., Bouferraa, Y., & Assi, H. I. (2022). Metabolic Syndrome: Updates on Pathophysiology and Management in 2021. *International Journal of Molecular Sciences*, 23(2), 786. <https://doi.org/10.3390/ijms23020786>
16. Ferrario, G., Baron, G., Gado, F., Della Vedova, L., Bombardelli, E., Carini, M., D'Amato, A., Aldini, G., & Altomare, A. (2022). Polyphenols from Thinned Young Apples: HPLC-HRMS Profile and Evaluation of Their Anti-Oxidant and Anti-Inflammatory Activities by Proteomic Studies. *Antioxidants*, 11(8), 1577. <https://doi.org/10.3390/antiox11081577>
17. Grima Serrano, A. (2010). El síndrome metabólico. *Acción Médica*.
18. Huang, P.-K., Lin, S.-R., Chang, C.-H., Tsai, M.-J., Lee, D.-N., & Weng, C.-F. (2019). Natural phenolic compounds potentiate hypoglycemia via inhibition of Dipeptidyl peptidase IV. *Scientific Reports*, 9(1). <https://doi.org/10.1038/s41598-019-52088-7>
19. Hussain, H., Green, I. R., Ali, I., Khan, I. A., Ali, Z., Al-Sadi, A. M., & Ahmed, I. (2017). Ursolic acid derivatives for pharmaceutical use: A patent review (2012-2016). *Expert Opinion on Therapeutic Patents*, 27(9), 1061-1072. <https://doi.org/10.1080/13543776.2017.1344219>

20. Jeong, G. H., Cho, J.-H., Park, E. K., & Kim, T. H. (2021). Enzymatic transformation products of phloretin as potent antiadipogenic compounds. *Bioscience, Biotechnology, and Biochemistry*, 85(12), 2352-2359. <https://doi.org/10.1093/bbb/zbab168>
21. Jiang, Y., Nan, H., Shi, N., Hao, W., Dong, J., & Chen, H. (2021). Chlorogenic acid inhibits proliferation in human hepatoma cells by suppressing noncanonical NF- κ B signaling pathway and triggering mitochondrial apoptosis. *Molecular Biology Reports*, 48(3), 2351-2364. <https://doi.org/10.1007/s11033-021-06267-3>
22. Jomova, K., Alomar, S. Y., Valko, R., Liska, J., Nepovimova, E., Kuca, K., & Valko, M. (2025). Flavonoids and their role in oxidative stress, inflammation, and human diseases. *Chemico-Biological Interactions*, 413, 111489. <https://doi.org/10.1016/j.cbi.2025.111489>
23. Khan, J., Deb, P. K., Priya, S., Medina, K. D., Devi, R., Walode, S. G., & Rudrapal, M. (2021). Dietary Flavonoids: Cardioprotective Potential with Antioxidant Effects and Their Pharmacokinetic, Toxicological and Therapeutic Concerns. *Molecules*, 26(13), 4021. <https://doi.org/10.3390/molecules26134021>
24. Kwon, E.-K., Lee, D.-Y., Lee, H., Kim, D.-O., Baek, N.-I., Kim, Y.-E., & Kim, H.-Y. (2010). Flavonoids from the Buds of *Rosa damascena* Inhibit the Activity of 3-Hydroxy-3-methylglutaryl-coenzyme A Reductase and Angiotensin I-Converting Enzyme. *Journal of Agricultural and Food Chemistry*, 58(2), 882-886. <https://doi.org/10.1021/jf903515f>
25. Libero, M. L., Lucarini, E., Recinella, L., Ciampi, C., Veschi, S., Piro, A., Chiavaroli, A., Acquaviva, A., Nilofar, N., Orlando, G., Generali, D., Ghelardini, C., Di Cesare Mannelli, L., Montero-Hidalgo, A. J., Luque, R. M., Ferrante, C., Menghini, L., Di Simone, S. C., Brunetti, L., & Leone, S. (2024). Anti-inflammatory and anti-

- hyperalgesic effects induced by an aqueous aged black garlic extract in rodent models of ulcerative colitis and colitis-associated visceral pain. *Phytotherapy Research*, 38(8), 4177-4188. <https://doi.org/10.1002/ptr.8270>
26. Liu, X., Fu, Y., Ma, Q., Yi, J., & Cai, S. (2021). Anti-Diabetic Effects of Different Phenolic-Rich Fractions from *Rhus Chinensis* Mill. *Fruits in vitro. eFood*, 2(1), 37-46. <https://doi.org/10.2991/efood.k.210222.002>
27. Maisto, M., Piccolo, V., Novellino, E., Schiano, E., Iannuzzo, F., Ciampaglia, R., Summa, V., & Tenore, G. C. (2022). Optimization of Phlorizin Extraction from Annurca Apple Tree Leaves Using Response Surface Methodology. *Antioxidants*, 11(10), 1933. <https://doi.org/10.3390/antiox11101933>
28. Martinez-Gonzalez, A. I., Díaz-Sánchez, Á. G., Rosa, L. A. D. L., Vargas-Requena, C. L., Bustos-Jaimes, I., & Alvarez-Parrilla, A. E. (2017). Polyphenolic Compounds and Digestive Enzymes: In Vitro Non-Covalent Interactions. *Molecules*, 22(4), 669. <https://doi.org/10.3390/molecules22040669>
29. Masenga, S. K., Kabwe, L. S., Chakulya, M., & Kirabo, A. (2023). Mechanisms of Oxidative Stress in Metabolic Syndrome. *International Journal of Molecular Sciences*, 24(9), 7898. <https://doi.org/10.3390/ijms24097898>
30. Medina-Torres, N., Ayora-Talavera, T., Espinosa-Andrews, H., Sánchez-Contreras, A., & Pacheco, N. (2017). Ultrasound Assisted Extraction for the Recovery of Phenolic Compounds from Vegetable Sources. *Agronomy*, 7(3), 47. <https://doi.org/10.3390/agronomy7030047>
31. Millán-Laleona, A., Cebollada, P., Caprioli, G., Piatti, D., Maggi, F., Pina, A., Gómez-Rincón, C., & López, V. (2025). Valorization of local regional apple (*Malus domestica* Borkh.) cultivars versus commercial samples from Spain: Phenolic compounds by HPLC-MS/MS, cytotoxicity and biological potential on nitric oxide radicals and

- lipoxygenase inhibition. *Journal of Functional Foods*, 124, 106631.
<https://doi.org/10.1016/j.jff.2024.106631>
32. Mustafa, A. M., Angeloni, S., Abouelenein, D., Acquaticci, L., Xiao, J., Sagratini, G., Maggi, F., Vittori, S., & Caprioli, G. (2022). A new HPLC-MS/MS method for the simultaneous determination of 36 polyphenols in blueberry, strawberry and their commercial products and determination of antioxidant activity. *Food Chemistry*, 367, 130743. <https://doi.org/10.1016/j.foodchem.2021.130743>
33. Nileeka Balasuriya, B. W., & Vasantha Rupasinghe, H. P. (2011). Plant flavonoids as angiotensin converting enzyme inhibitors in regulation of hypertension. *Functional Foods in Health and Disease*, 1(5), 172. <https://doi.org/10.31989/ffhd.v1i5.132>
34. Nirmal, N., Khanashyam, A., Mundanat, A., Shah, K., Babu, K., Thorakkattu, P., Al-Asmari, F., & Pandiselvam, R. (2023). Valorization of Fruit Waste for Bioactive Compounds and Their Applications in the Food Industry. *Foods*, 12(3), 556. <https://doi.org/10.3390/foods12030556>
35. Noubiap, J. J., Nansseu, J. R., Lontchi-Yimagou, E., Nkeck, J. R., Nyaga, U. F., Ngouo, A. T., Tounouga, D. N., Tianyi, F.-L., Foka, A. J., Ndoadoumgué, A. L., & Bigna, J. J. (2022). Geographic distribution of metabolic syndrome and its components in the general adult population: A meta-analysis of global data from 28 million individuals. *Diabetes Research and Clinical Practice*, 188, 109924. <https://doi.org/10.1016/j.diabres.2022.109924>
36. Nunes, A. R., Alves, G., Falcão, A., Lopes, J. A., & Silva, L. R. (2025). Phenolic Acids from Fruit By-Products as Therapeutic Agents for Metabolic Syndrome: A Review. *International Journal of Molecular Sciences*, 26(8), 3834. <https://doi.org/10.3390/ijms26083834>

37. Ojha, S., Islam, B., Charu, C., Adem, A., & Aburawi, E. (2015). Insight into the mechanism of polyphenols on the activity of HMGR by molecular docking. *Drug Design, Development and Therapy*, 4943. <https://doi.org/10.2147/dddt.s86705>
38. Pang, Z., Chong, J., Zhou, G., de Lima Morais, D. A., Chang, L., Barrette, M., Gauthier, C., Jacques, P.-É., Li, S., & Xia, J. (2021). MetaboAnalyst 5.0: Narrowing the gap between raw spectra and functional insights. *Nucleic Acids Research*, 49(W1), W388-W396. <https://doi.org/10.1093/nar/gkab382>
39. Popiolek-Kalisz, J., & Glibowski, P. (2023). Apple Peel Supplementation Potential in Metabolic Syndrome Prevention. *Life*, 13(3), 753. <https://doi.org/10.3390/life13030753>
40. Procházková, D., Boušová, I., & Wilhelmová, N. (2011). Antioxidant and prooxidant properties of flavonoids. *Fitoterapia*, 82(4), 513-523. <https://doi.org/10.1016/j.fitote.2011.01.018>
41. Ran, X., Zhang, M., Wang, Y., & Adhikari, B. (2019). Novel technologies applied for recovery and value addition of high value compounds from plant byproducts: A review. *Critical Reviews in Food Science and Nutrition*, 59(3), 450-461. <https://doi.org/10.1080/10408398.2017.1377149>
42. Sahu, S. C., Zheng, J., Graham, L., Chen, L., Ihrle, J., Yourick, J. J., & Sprando, R. L. (2014). Comparative cytotoxicity of nanosilver in human liver HepG2 and colon Caco2 cells in culture. *Journal of Applied Toxicology*, 34(11), 1155-1166. <https://doi.org/10.1002/jat.2994>
43. Sanz-Buenhombre, M., Villanueva, S., Moro, C., Tomás-Cobos, L., Viadel, B., & Guadarrama, A. (2016). Bioavailability and the mechanism of action of a grape extract rich in polyphenols in cholesterol homeostasis. *Journal of Functional Foods*, 21, 178-185. <https://doi.org/10.1016/j.jff.2015.11.044>

44. Tian, C., Hao, L., Yi, W., Ding, S., & Xu, F. (2020). Polyphenols, Oxidative Stress, and Metabolic Syndrome. *Oxidative Medicine and Cellular Longevity*, 2020, 1-2. <https://doi.org/10.1155/2020/7398453>
45. Wei, M., Wang, H., Ma, T., Ge, Q., Fang, Y., & Sun, X. (2021). Comprehensive Utilization of Thinned Unripe Fruits from Horticultural Crops. *Foods*, 10(9), 2043. <https://doi.org/10.3390/foods10092043>
46. Yang, C. S., Landau, J. M., Huang, M.-T., & Newmark, H. L. (2001). INHIBITION OF CARCINOGENESIS BY DIETARY POLYPHENOLIC COMPOUNDS. *Annual Review of Nutrition*, 21(1), 381-406. <https://doi.org/10.1146/annurev.nutr.21.1.381>
47. Yeh, W.-J., Hsia, S.-M., Lee, W.-H., & Wu, C.-H. (2017). Polyphenols with antiglycation activity and mechanisms of action: A review of recent findings. *Journal of Food and Drug Analysis*, 25(1), 84-92. <https://doi.org/10.1016/j.jfda.2016.10.017>
48. Yoshida, K., Zhao, P., Zhang, L., Abernethy, D. R., Rekić, D., Reynolds, K. S., Galetin, A., & Huang, S.-M. (2017). In Vitro–In Vivo Extrapolation of Metabolism- and Transporter-Mediated Drug–Drug Interactions—Overview of Basic Prediction Methods. *Journal of Pharmaceutical Sciences*, 106(9), 2209-2213. <https://doi.org/10.1016/j.xphs.2017.04.045>
49. Zheng, H.-Z., Cui, C.-L., Jeong, W.-S., & Chung, S.-K. (2020). Anti-inflammatory effect of unripe apple polyphenols-chitoooligosaccharides microcapsule against LPS-induced RAW 264.7 cells. *Applied Biological Chemistry*, 63(1), 51. <https://doi.org/10.1186/s13765-020-00533-x>
50. Zheng, H.-Z., Kim, Y.-I., & Chung, S.-K. (2012). A profile of physicochemical and antioxidant changes during fruit growth for the utilisation of unripe apples. *Food Chemistry*, 131(1), 106-110. <https://doi.org/10.1016/j.foodchem.2011.08.038>

Table 1. Quantitative determination of polyphenols in the thinned apple extracts (mg of phenolic compounds kg⁻¹ of extract) by using HPLC-MS/MS triple quadrupole.

Compounds	Amarilla de Octubre	Pomera de Pomes Agrias	Camuesa de Castellano Daroca	Daroca-02	Vadiello-02	Verde Doncella	Reineta	Golden Smoothee	Fuji	Royal Gala
Hydroxybenzoic Acids										
Gallic acid	23.11	11.60	n.d.	6.78	n.d.	12.43	2.42	n.d.	n.d.	n.d.
3-Hydroxy benzoic acid	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
4-Hydroxy benzoic acid	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Cinnamic Acids										
Chlorogenic acid	1945.66	1144.56	910.43	567.98	234.97	414.87	210.98	111.87	812.85	312.85
Neochlorogenic acid	423.56	34.80	5.98	n.d.	n.d.	n.d.	111.34	45.90	60.98	n.d.
Caffeic acid	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Vanillic acid	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Syringic acid	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
P-Coumaric acid	13.44	4.05	n.d.	n.d.	n.d.	22.81	n.d.	n.d.	11.67	13.98
Ferulic acid	n.d.	n.d.	5.67	3.97	4.9	n.d.	1.98	n.d.	n.d.	n.d.
Ellagic acid	144.13	44.86	34.85	45.08	34.87	199.54	n.d.	109.98	48.80	106.86
Trans-cinnamic acid	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Stilbenes										
Resveratrol	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Dihydrochalcones										
Phlorizin	323.67	132.98	30.59	134.80	76.98	138.90	134.97	198.87	207.43	107.32
Phloretin	0.97	1.43	0.56	0.16	n.d.	2.22	1.14	0.34	1.14	1.09
Flavonols										
Quercetin	45.98	87.82	66.6	55.53	33.34	9.98	33.07	45.97	113.07	98.55
Kaempferol	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Myricetin	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Isorhamnetin	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.

Table 1 (continued)

Kaempferol-3-glucoside	28.98	11.97	8.9	11.37	33.85	14.98	18.71	23.33	45.66	66.75
Quercitrin	1567.98	1447.40	442.78	456.86	954.97	45.89	342.78	517.20	824.56	723.97
Rutin	698.09	442.78	732.78	540.08	1142.78	923.98	122.78	322.78	1134.33	934.99
Hyperoside	1945.87	936.44	2345.28	1237.30	636.43	4389.09	1236.43	1123.89	2709.70	3809.05
Isoquercitrin	1031.45	2134.08	1678.34	2224.90	1228.09	3550.87	3755.56	4770.77	1704.55	1199.82
Flavan-3-ols										
Catechin	502.54	142.78	456.96	135.96	452.08	248.99	110.09	98.43	111.65	232.07
Procyanidin A2	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	4.76	5.98
Procyanidin B2	1520.56	942.78	644.11	1441.87	742.19	1456.08	1232.78	992.42	986.9	776.09
Epicatechin	979.45	535.87	1077.12	1255.01	996.31	766.12	323.56	345.67	1412.45	1186.63
Flavanones										
Naringenin	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Hesperidin	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Anthocyanins										
Delphinidin-3,5-diglucoside	1960.71	1613.55	1777.93	1223.56	1533.45	1134.08	868.56	999.03	3567.45	4048.11
Delphinidin-3-galactoside	11.11	17.77	n.d.	n.d.	n.d.	7.06	18.8	11.08	249.87	153.43
Cyanidin-3-glucoside	n.d.	n.d.	n.d.	n.d.	n.d.	0.11	n.d.	n.d.	3.11	2.45
Petunidin-3-glucoside	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Pelargonidin-3-glucoside	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Pelargonidin-3-rutinoside	n.d.	n.d.	n.d.	n.d.	n.d.	1.34	n.d.	n.d.	n.d.	n.d.
Malvidin-3-galactoside	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Total Flavonoid Content	10617.36	8447.65	9261.95	8717.40	7830.47	12689.69	8199.23	9449.78	13076.63	13346.30
Total Phenolic Content	13167.26	9687.52	10218.88	9341.21	8105.21	13339.34	8525.95	9717.53	14010.93	13779.99

n.d.: not detected




Fig 1. Chemical structures of the predominant phenolic compounds in thinned apple by-products




Fig 2. Hierarchical clustering heatmap analysis. Heatmap values are Z-score normalized by compound. Red cells ($Z > 0$) indicate samples enriched in a given phenolic relative to its dataset mean, enabling row-wise comparison of enrichment patterns.




Fig 3. PCA score plot of phenolic groups




Fig 4. Cell viability (%) determined in HepG2 (A), Caco-2 (B) and HFF-1 (C) cells treated with thinned apple extracts at concentrations ranging from 15.62 to 500 $\mu\text{g mL}^{-1}$. Results are presented as mean \pm SEM from at least three independent experiments.

Table 2. Results of antidiabetic, antiglycation, antioxidant activity hypolipidemic, hypocholesterolemic and hypotensive activity against α -glucosidase, α -amylase, DPP-4, AGEs formation, radical superoxid, lipase, HMG-COA reductase and ACE. Data are expressed as mean \pm SEM from at least three independent experiments. IC₅₀ values were calculated using nonlinear regression. Statistical significance: ns = not significant; *p < 0.05; **p < 0.001; ***p < 0.0005; ****p < 0.0001 vs. control

IC ₅₀ values ($\mu\text{g mL}^{-1}$) of the in vitro assays								
Extracts	α -Glucosidase	α -Amylase	DPP-4	AGEs	O ₂ ^{•-} scavenging	Lipase	HMGR	ACE
Verde Doncella	361.56 \pm 6.04 ^{ns}	75.84 \pm 72.132****	766.89 \pm 29.57****	203.44 \pm 5.61****	2.75 \pm 0.26 ^{ns}	363.90 \pm 25.91***	100.88 \pm 3.93****	42.17 \pm 3.11 ^{ns}
Golden Smoothee	1342.37 \pm 58.26****	n.a	114.76 \pm 49.74****	381.16 \pm 24.83****	4.03 \pm 0.48 ^{ns}	875.02 \pm 13.57****	469.49 \pm 8.04****	209.28 \pm 17.26****
Reineta	486.47 \pm 8.85*	174.71 \pm 69.02****	193.10 \pm 48.02****	189.36 \pm 13.15****	2.47 \pm 0.50 ^{ns}	225.43 \pm 6.31*	149.39 \pm 9.39****	84.34 \pm 11.58*
Fuji	826.19 \pm 51.25****	n.a	281.03 \pm 133.44****	434.48 \pm 15.64****	2.13 \pm 0.29 ^{ns}	397.1 \pm 44.77****	n.a	206.28 \pm 12.12****
Royal Gala	1644.86 \pm 43.83****	n.a	1763 \pm 89.71****	419.44 \pm 20.90****	4.18 \pm 0.46 ^{ns}	635.65 \pm 13.48****	n.a	290.02 \pm 22.81****
Amarilla de Octubre	105.60 \pm 7.23 ^{ns}	06.94 \pm 17.23****	628.03 \pm 21.09***	49.77 \pm 3.52 ^{ns}	1.18 \pm 0.21 ^{ns}	419.31 \pm 17.76****	n.a	55.94 \pm 1.18 ^{ns}
Pomera de Pomes Agrias	123.71 \pm 11.70 ^{ns}	337.12 \pm 59.73*	480.63 \pm 3.39*	60.08 \pm 3.01 ^{ns}	1.98 \pm 0.02 ^{ns}	447.02 \pm 25.45****	n.a	81.32 \pm 10.33*
Vadiello-02	244.14 \pm 15.08 ^{ns}	194.79 \pm 22.88 ^{ns}	110.34 \pm 66.72****	871.90 \pm 9.20****	9.74 \pm 2.78*	948.38 \pm 5.26****	233.04 \pm 17.78****	287.24 \pm 21.32****
Daroca-02	1484.60 \pm 66.79****	n.a	496.99 \pm 131.97****	343.74 \pm 14.95****	4.10 \pm 0.92 ^{ns}	1332.2 \pm 83.88****	n.a	90.91 \pm 12.09*
C. de Castellano de Daroca	1733.29 \pm 112.18****	n.a	617.74 \pm 116.73****	835.12 \pm 34.16****	7.12 \pm 0.45 ^{ns}	901.74 \pm 89.56****	n.a	242.44 \pm 20.55****
Acarbose	220 \pm 64.10	7.72 \pm 1.89	-	-	-	-	-	-
Sitagliptin	-	-	0.04 \pm 0.01	-	-	-	-	-
Aminoguanidine	-	-	-	55.01 \pm 5.63	-	-	-	-
Quercetin	-	-	-	-	4.16 \pm 0.24	-	-	-
Orlistat	-	-	-	-	-	32.20 \pm 4.28	-	-
Pravastatin	-	-	-	-	-	-	0.012 \pm 0.0001	-

Captopril

-

-

-

-

-

-

-

 0.003 ± 0.001

n.a: no activity

Journal Pre-proof




Fig 5. Pearson correlation heatmap showing the relationships between IC_{50} values of the enzymes obtained from all in vitro bioassays and the individual phenolic content. Each cell displays the Pearson correlation coefficient, where a value of 1.0 denotes a positive linear correlation (red) and -1.0 indicates negative correlation (blue).

Highlights

- Thinned apples are a common by-product produced in abundant quantities
- Hyperoside, isoquercitrin, and delphinidin-3,5-diglucoside emerged as the main phenolic compounds
- Autochthonous and commercial cultivars exhibited bioactivity
- Thinned apples can be used as sources for the development of nutraceuticals and bioactive ingredients

Journal Pre-proof