

Draft Genome Sequence of Two Strains of *Xanthomonas arboricola* Isolated from *Prunus persica* Which Are Dissimilar to Strains That Cause Bacterial Spot Disease on *Prunus* spp.

Jerson Garita-Cambronero,^a Ana Palacio-Bielsa,^b María M. López,^c  Jaime Cubero^a

Departamento de Protección Vegetal, Instituto Nacional de Investigación y Tecnología Agraria, Madrid, Spain^a; Centro de Investigación y Tecnología Agroalimentaria de Aragón, Instituto Agroalimentario de Aragón-IA2 (CITA-Universidad de Zaragoza), Zaragoza, Spain^b; Instituto Valenciano de Investigaciones Agrarias, Valencia, Spain^c

The draft genome sequences of two strains of *Xanthomonas arboricola*, isolated from asymptomatic peach trees in Spain, are reported here. These strains are avirulent and do not belong to the same phylogroup as *X. arboricola* pv. *pruni*, a causal agent of bacterial spot disease of stone fruits and almonds.

Received 18 July 2016 Accepted 19 July 2016 Published 8 September 2016

Citation Garita-Cambronero J, Palacio-Bielsa A, López MM, Cubero J. 2016. Draft genome sequence of two strains of *Xanthomonas arboricola* isolated from *Prunus persica* which are dissimilar to strains that cause bacterial spot disease on *Prunus* spp. *Genome Announc* 4(5):e00974-16. doi:10.1128/genomeA.00974-16.

Copyright © 2016 Garita-Cambronero et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Jaime Cubero, cubero@inia.es.

Xanthomonas arboricola is a Gram-negative species that comprises nine subspecific phylogroups (1, 2). In addition, some strains that differ from the pathovars *X. arboricola* pv. *pruni* and *X. arboricola* pv. *juglandis*, isolated from the plant genera *Prunus* and *Juglans*, have also been described. These strains are unable to cause disease because they lack some essential characteristics related to pathogenesis, such as the type III secretion system and related effectors (3, 4).

The two new genomes presented here, which belong to the avirulent strains CITA 14 and CITA 124, were compared to the genomes of other *X. arboricola* strains (3–9) and could be helpful in understanding the pathogenesis and evolution in this species. Moreover, the genomes will be useful for generating accurate diagnostic tools to differentiate the quarantine pathogen *X. arboricola* pv. *pruni* from other nontarget strains, thereby avoiding unnecessary control measures that could result in high economic losses as occurred with other xanthomonads (10, 11).

Strains CITA 14 and CITA 124 were isolated from asymptomatic peach trees (*Prunus persica*) in Spain and were deposited in the Centro de Investigación y Tecnología Agroalimentaria de Aragón (CITA) in Zaragoza, Spain. To confirm their identification, a multilocus sequence analysis using partial sequences of the housekeeping genes *dnaK*, *fyuA*, *gyrB*, and *rpoD* was performed (12). By means of such analysis, these strains were not clustered into any of the established pathovars of *X. arboricola*. CITA 14 and CITA 124 were enclosed in a group with CITA 44, another atypical strain of *X. arboricola* isolated from the rootstock Santa Lucía SL-64 (*Prunus mahaleb*) (4), which shows sequence identities of 98.3% with strain CITA 14 and 98.9% with strain CITA 124.

Genome sequencing was performed using Ion Torrent sequencing technology (PGM, Life Technologies) at STAB VIDA, Caparica, Portugal (733.06 Mb, 100-fold coverage for CITA 14 and 403.99 Mb, 50-fold coverage for CITA 124). *De novo* assembly was conducted using CLC Genomics Workbench version 8.5.1 (CLC bio, Denmark) and MIRA version 4.0 (13), and the subsequent merging of the two assemblies was done using Sequencher

version 5.4 (Gene Codes Corporation, USA). Total lengths of 4,746,212 bp (65.60% G+C content) and 4,752,241 bp (65.8% G+C content) were generated and placed in 72 (N_{50} , 108,424 bp; maximum contig length, 265,354 bp) and 128 contigs (N_{50} , 60,818 bp; maximum contig length 189,085) for CITA 14 and CITA 124, respectively.

Annotations were conducted using the NCBI Prokaryotic Genome Annotation Pipeline version 3.1 (14) and the Rapid Annotations using Subsystem Technology server (15). The annotation for strain CITA 14 detected 3,974 coding sequences, four rRNAs, and 53 tRNAs, representing 439 subsystems; meanwhile, for strain CITA 124, 3,797 coding sequences, three rRNAs, and 50 tRNAs, representing 442 subsystems, were detected. A detailed genome comparative analysis will be addressed in a future publication.

Accession number(s). These whole-genome shotgun projects have been deposited in DDBJ/EMBL/GenBank under the accession numbers [LXIB00000000](https://accession.gtrdb.org/acc/show/LXIB00000000) for strain CITA 14 and [LXKK00000000](https://accession.gtrdb.org/acc/show/LXKK00000000) for strain CITA 124. The versions described in this paper are the first versions, LXIB01000000 and LXKK01000000.

ACKNOWLEDGMENTS

This work was supported financially by the Instituto Nacional de Investigación y Tecnología Agraria y Alimentaria (INIA) project RTA2014-00018-CO2-01. J. Garita-Cambronero held a Ph.D. fellowship from the Spanish Government (Ministerio de Educación, Cultura y Deporte; fellowship FPU12/01000).

FUNDING INFORMATION

This work, including the efforts of Jerson Garita-Cambronero, was funded by Ministerio de Educación, Cultura y Deporte (MECD) (FPU12/01000). This work was funded by Instituto Nacional de Investigación y Tecnología Agraria y Alimentaria (INIA) (RTA2014-00018-CO2-01).

REFERENCES

- Vauterin L, Hoste B, Kersters K, Swings J. 1995. Reclassification of *Xanthomonas*. *Int J Syst Bacteriol* 45:472–489. <http://dx.doi.org/10.1099/00207713-45-3-472>.

2. Fischer-Le Saux M, Bonneau S, Essakhi S, Manceau C, Jacques M-A. 2015. Aggressive emerging pathovars of *Xanthomonas arboricola* represent widespread epidemic clones distinct from poorly pathogenic strains, as revealed by multilocus sequence typing. *Appl Environ Microbiol* 81: 4651–4668. <http://dx.doi.org/10.1128/AEM.00050-15>.
3. Garita-Cambronero J, Palacio-Bielsa A, López MM, Cubero J. 2016. Draft genome sequence for virulent and avirulent strains of *Xanthomonas arboricola* isolated from *Prunus* spp. in Spain. *Stand Genomic Sci* 11:12. <http://dx.doi.org/10.1186/s40793-016-0132-3>.
4. Cesbron S, Briand M, Essakhi S, Gironde S, Boureau T, Manceau C, Fischer-Le Saux M, Jacques M-A. 2015. Comparative genomics of pathogenic and nonpathogenic strains of *Xanthomonas arboricola* unveil molecular and evolutionary events linked to pathoadaptation. *Front Plant Sci* 6:1126. <http://dx.doi.org/10.3389/fpls.2015.01126>.
5. Garita-Cambronero J, Sena-Vélez M, Palacio-Bielsa A, Cubero J. 2014. Draft genome sequence of *Xanthomonas arboricola* pv. pruni strain Xap33, causal agent of bacterial spot disease on almond. *Genome Announc* 2(3): e00440-14. <http://dx.doi.org/10.1128/genomeA.00440-14>.
6. Ibarra Caballero J, Zerillo MM, Snelling J, Boucher C, Tisserat N. 2013. Genome sequence of *Xanthomonas arboricola* pv. corylina, isolated from Turkish filbert in Colorado. *Genome Announc* 1(3):e00246-13. <http://dx.doi.org/10.1128/genomeA.00246-13>.
7. Harrison J, Grant MR, Studholme DJ. 2016. Draft genome sequences of two strains of *Xanthomonas arboricola* pv. celebensis isolated from banana plants. *Genome Announc* 4(1):e01705-15. <http://dx.doi.org/10.1128/genomeA.01705-15>.
8. Higuera G, González-Escalona N, Véliz C, Vera F, Romero J. 2015. Draft genome sequences of four *Xanthomonas arboricola* pv. juglandis strains associated with walnut blight in Chile. *Genome Announc* 3(5): e01160-15. <http://dx.doi.org/10.1128/genomeA.01160-15>.
9. Ignatov AN, Kyrova EI, Vinogradova SV, Kamionskaya AM, Schaad NW, Luster DG. 2015. Draft genome sequence of *Xanthomonas arboricola* strain 3004, a causal agent of bacterial disease on barley. *Genome Announc* 3(1):e01572-14. <http://dx.doi.org/10.1128/genomeA.01572-14>.
10. Jalan N, Aritua V, Kumar D, Yu F, Jones JB, Graham JH, Setubal JC, Wang N. 2011. Comparative genomic analysis of *Xanthomonas axonopodis* pv. citrumelo F1, which causes citrus bacterial spot disease, and related strains provides insights into virulence and host specificity. *J Bacteriol* 193:6342–6357. <http://dx.doi.org/10.1128/JB.05777-11>.
11. Graham JH, Gottwald TR. 1991. Research perspectives on eradication of citrus bacterial diseases in Florida. *Plant Dis* 75:1193–1200. <http://dx.doi.org/10.1094/PD-75-1193>.
12. Young JM, Park DC, Shearman HM, Fargier E. 2008. A multilocus sequence analysis of the genus *Xanthomonas*. *Syst Appl Microbiol* 31: 366–377. <http://dx.doi.org/10.1016/j.syapm.2008.06.004>.
13. Chevreux B, Wetter T, Suhai S. 1999. Genome sequence assembly using trace signals and additional sequence information, p. 45–56. *In* Wingender E (ed), *Computer science and biology: proceedings of the German Conference on Bioinformatics (GCB) 1999*, Hannover, Germany. GBF-Braunschweig, Department of Bioinformatics, Braunschweig, Germany.
14. Angiuoli SV, Gussman A, Klimke W, Cochrane G, Field D, Garrity G, Kodira CD, Kyrpides N, Madupu R, Markowitz V, Tatusova T, Thomson N, White O. 2008. Toward an online repository of standard operating procedures (SOPs) for (meta)genomic annotation. *Omic* 12:137–141. <http://dx.doi.org/10.1089/omi.2008.0017>.
15. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <http://dx.doi.org/10.1186/1471-2164-9-75>.