Session 41

Theatre 8

Relationship between resource use, efficiency and sustainability of sheep-crop farming systems

T. Rodríguez-Ortega^{1,2}, A. Bernués^{1,2} and A. Olaizola^{1,3}

¹Instituto Agroalimentario de Aragón – IA2 (CITA-Universidad de Zaragoza), Av. Montañana 930, 50059 Zaragoza, Spain, ²Centro de Investigación y Tecnología Agroalimentaria de Aragón (CITA), Av. Montañana 930, 50059 Zaragoza, Spain, ³Universidad de Zaragoza, Departamento de Ciencias Agrarias y del Medio Natural, C/ Miguel Servet 177, 50013 Zaragoza, Spain; abernues@aragon.es

Farming systems determine the resource embeddedness in agricultural products. Quantification of efficiency and sustainability is challenging in Mediterranean sheep-crop farming systems due to their different specialization, crop-animal integration and intensification. Emergy analysis computes the different qualities of energies involved in a production process and expresses them in equivalent solar energy. It provides emergy-based indicators such as intensity (emergy used per hectare and year), efficiency (emergy invested per energy obtained), and sustainability (ratio of self-sufficiency to environmental loading). Our objective was to evaluate these indicators on three representative farms of Mediterranean sheep-crop farming systems of Aragón (Spain) at two different levels: the product and the farm level (including lamb meat, permanent crops, and rainfed and irrigated arable crops). Our results show that the pasture-based sheep farm had the lowest intensity and efficiency and the highest sustainability, as opposite to the partially-integrated farm, while the fully-integrated farm obtained intermediate scores. Sheep products were less intensive and more sustainable than crops due to their capacity to use renewable natural resources (i.e. pastures). Since emergy increases with the hierarchical level of the food chain (e.g. from crops to meat), efficiency should be compared between products of similar trophic level. Lamb meat production was 1.9 and 1.3 times more intensive and efficient, respectively, in the partially-integrated farm than in the pasture-based sheep farm, but it was 5.1 times less sustainable. Increasing the boundaries of emergy analysis from the farm to the regional level could show strategies of resource sharing among farms for increased regional sustainability.

Book of Abstracts of the 69th Annual Meeting of the European Federation of Animal Science

Dubrovnik, Croatia, 27th – 31st August, 2018



EAAP Scientific Committee:

E. Strandberg G. Savoini H.A.M. Spoolder H. Sauerwein M. Tichit M. Klopčič J. Conington E.F. Knol A.S. Santos T. Veldkamp I. Halachmi G.E. Pollott



publication may be translated, reproduced, stored in a computerised system or published in any form or in any manner, including electronic, mechanical, reprographic or photographic, without prior written permission from the publisher: Wageningen Academic Publishers P.O. Box 220 6700 AE Wageningen The Netherlands www.WageningenAcademic.com copyright@WageningenAcademic.com

This work is subject to copyright. All rights are reserved, whether the whole or part of the material is concerned. Nothing from this

The individual contributions in this publication and any liabilities arising from them remain the responsibility of the authors.

The designations employed and the presentation of material in this publication do not imply the expression of any opinion whatsoever on the part of the European Federation of Animal Science concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries.

The publisher is not responsible for possible damages, which could be a result of content derived from this publication.

EAN: 9789086863235 e-EAN: 9789086868711 ISBN: 978-90-8686-323-5 e-ISBN: 978-90-8686-871-1 DOI: 10.3920/978-90-8686-871-1

ISSN 1382-6077

First published, 2018

© Wageningen Academic Publishers The Netherlands, 2018



Wageningen Academic Publishers