

EWRS Working Group "Weed Management Systems in Vegetables"



Weeds and weed management in lettuce

F. Tei*, A. Cirujeda, A. Dobrzański, R. J. Froud-Williams, H. Mennan, R. Neuweiler, E. Pannacci, J. Qasem, F. Rocha, A. Simončič, A. Verschwele, R. van der Weide, C. Zaragoza

* Dept. Agricultural and Environmental Sciences, University of Perugia, Italy - f.tei@unipg.it

Objective

Information about key weeds, new weeds or species that have recently become problematic, effect of competition, weed management programmes in integrated and organic production, approved herbicides and those currently undergoing registration for use in lettuce grown in Germany (D), Italy (I), Jordan (HKJ), The Netherlands (NL), Poland (PL), Portugal (P), Slovenia (SLO), Spain (E), Switzerland (CH), Turkey (TR) and United Kingdom (UK) was collected.

Lettuces in surveyed countries

Table with 2 columns: Country, ha. Lists countries like Spain, Turkey, Italy, Germany, etc. with their respective lettuce production in hectares.

Although the majority of the lettuce crop is grown as an outdoor spring-summer crop (see table above), early and late productions are obtained by non-woven materials (fleece), clear/white plastic mulch, row covers and greenhouse cultivations to satisfy year-round consumption.

Weeds

Key weeds

Weeds that are most problematic in lettuce are those that are taxonomically related and not controlled by herbicides, in particular, Galinsoga spp., Anthemis spp., Matricaria spp., Senecio vulgaris and Sonchus oleraceus.

Weeds are becoming important

Rorippa sylvestris in CH. Abutilon theophrasti, Galinsoga parviflora, Xanthium spinosum, X. strumarium in P. Aethusa cynapium, Bidens tripartita in D. Cruciferae in SLO. Oxalis pes-caprae and Calendula arvensis in I.

Weed competition


Transplanted crops rarely suffer severe weed competition, but late maturing varieties and all drilled crops are particularly susceptible due to low initial growth rates. Typically there is no critical period of competition such that a single weeding 2-3 weeks after transplanting is sufficient to prevent yield loss.

Approved active ingredients

Table with columns: Active ingredients, Application time, and registration status in various countries (UK, PL, NL, D, CH, SLO, I, E, P, HKJ).

(1) pres = pre-sowing; pree = pre-emergence; pret = pre-transplanting; poste = post-emergence; postt = post-transplanting

No pre- or post-emergence/transplanting herbicides are registered in Turkey



Conventional weed control


a pre-emergence or pre-transplanting treatment + (if necessary) a post-emergence / post-transplanting treatment

Integrated Weed Management System

- 1. false seedbed technique followed by shallow harrowings or by glyphosate or glufosinate-ammonium application
2. transplanting
3. pre-transplanting herbicide application or black plastic mulching
4. post-transplanting inter-row hoeing

Organic production

- 1. false seedbed technique followed by shallow harrowing
2. transplanting
3. split-hoeing and/or finger weeding or starch-based biodegradable mulching
4. hand-weeding



Weeds and weed management in lettuce

F. Tei, A. Cirujeda, A. Dobrzański, R. J. Froud-Williams, H. Mennan, R. Neuweiler, E. Pannacci, J. Qasem, F. Rocha, A. Simončič, A. Verschwele, R. van der Weide, C. Zaragoza
Dept. of Agricultural and Environmental Sciences, University of Perugia, Italy
f.tei@unipg.it

The EWRS Working Group "Weed Management Systems in Vegetables" was established with the aim of collecting and disseminating information and results on weeds and weed control strategies in vegetables, identifying gaps in knowledge and defining new research projects. So far the working group has published reviews on onions, tomatoes, carrots, peas and cabbages. Information about key weeds, new weeds or species that have recently become problematic, effect of competition, weed management programmes in integrated and organic production, approved herbicides and those currently undergoing registration for use in lettuce grown in Germany (D), Italy (I), Jordan (HKJ), The Netherlands (NL), Poland (PL), Portugal (P), Slovenia (SLO), Spain (E), Switzerland (CH), Turkey (TR) and United Kingdom (UK) was collected.

Although the majority of the lettuce crops is grown as an outdoor spring-summer crop (E 37000 ha, TR 20000 ha, I 18000 ha, D 12500 ha, UK 7000 ha, HKJ 1900 ha, CH 1700 ha, NL 1300 ha, PL 600 ha, SLO 350 ha,), early and late productions are obtained by non-woven materials (fleece), clear/white plastic mulch, row covers and greenhouses cultivations for satisfying a year-round consumption. Black polyethylene and black biodegradable plastic (Mater-Bi) mulching are usually preferred where weed control is a priority in comparison to yield earliness. Lettuces are mainly transplanted as modules but may be also direct-drilled. The weed communities are commonly very species rich and their composition is highly variable in relation to climate, soil and crop period.

Weeds that are most problematic in lettuce are those that are taxonomically related and not controlled by herbicides, in particular, *Galinsoga* spp., *Anthemis* spp., *Matricaria* spp., *Senecio vulgaris* and *Sonchus oleraceus*.

However, *Capsella bursa-pastoris*, *Diplotaxis* spp., *Fumaria officinalis*, *Papaver rhoeas*, *Veronica* spp., *Viola arvensis*, *Stellaria media*, *Urtica urens*, *Chenopodium* spp. and *Poa annua* may also be key weeds in early cultivations whilst *Amaranthus* spp., *Polygonum* spp., *Portulaca oleracea*, *Digitaria sanguinalis* and *Echinochloa crus-galli* in late cultivations.

Some species are becoming important: *Rorippa sylvestris* in CH, *Abutilon theophrasti*, *Galinsoga parviflora*, *Xanthium spinosum*, *X. strumarium* in P, *Aethusa cynapium*, *Bidens tripartita* in D, *Cruciferae* in SLO, and *Oxalis pes-caprae* and *Calendula arvensis* in I.

Transplanted crops rarely suffer severe weed competition, but late maturing varieties and all drilled crops are particularly susceptible due to low initial growth rates. Typically there is no critical period of competition such that a single weeding 2-3 weeks after transplanting is sufficient to prevent yield loss. However, there is a zero tolerance of weeds that may hinder hand-harvesting or lead to contamination of minimally processed lettuces.

Conventional weed control is based on herbicide application; options are restricted to trifluralin (I, UK, HKJ), chlorpropham (I, UK, NL), pendimethalin (E, I, UK, D, CH, HKJ), propachlor (I, UK), propyzamide (E, I, UK, D, CH, P, PL, SLO), carbetamide (NL), benfluralin (E, I), chlorthal (E, I), oxadiargil (E), oxadiazon (I), oxyfluorfen (HKJ) and graminicides (E, I, P, D, SLO, HKJ). No pre- or post-emergence/transplanting herbicides are registered in TR. In CH, flufenazet 0.6 L/ha applied immediately before planting showed a total control of *Galinsoga* spp. Common strategy for organic production is a false seedbed technique, mechanical (i.e. split-hoeing and/or finger weeding) and manual weeding; planting on ridges and starch-based biodegradable mulching are increasing.

Further and more detailed information can be found in the WG web site at www.veg.ewrs.org