



## Evolution of pastoral livestock farming on arid rangelands in the last 15 years



H. Rjili <sup>a,b,\*</sup>, E. Muñoz-Ulecia <sup>c,d</sup>, A. Bernués <sup>c,d</sup>, M. Jaouad <sup>a</sup>, D. Martin-Collado <sup>c,d</sup>

<sup>a</sup> Laboratory of Economy and Rural Societies, Arid Regions Institute of Medenine (IRA), Gabes University, 4119 Medenine, Tunisia

<sup>b</sup> National Agronomic Institute of Tunis (INAT), University of Carthage, 1082 Tunis, Tunisia

<sup>c</sup> Departamento de Ciencia Animal, Centro de Investigación y Tecnología Agroalimentaria de Aragón (CITA), Avda. Montañana 930, 50059 Zaragoza, Spain

<sup>d</sup> Instituto Agroalimentario de Aragón – IA2 (CITA-Universidad de Zaragoza), Zaragoza, Spain

### ARTICLE INFO

#### Article history:

Received 17 September 2021

Revised 8 February 2023

Accepted 10 February 2023

Available online 24 February 2023

#### Keywords:

Camels

Farm dynamics

Farm typologies

Sheep

Temporal dynamics

### ABSTRACT

Livestock farming in arid rangelands constitutes a key component in the agricultural sector, particularly in developing countries. Farms have rapidly changed in recent decades, which has resulted in the modification of their structure, management and economic performance. Nowadays, livestock production in arid rangelands is threatened by climate change, coupled with the impact of complex interactions among social, economic and political factors. The present study analyses the main changes that have occurred on farms in the arid rangelands of south Tunisia from 2004 to 2019 and discusses the factors that explain the geographical patterns of such changes. Data were collected through face-to-face questionnaires with 73 farmers in two years (2004 and 2019). Information included farm structure and management, resources use and economic performance. Multivariate statistical methods analysed the differences in farm typologies between dates and the different pathways of change. Results showed that most farms increased herd size and cereal area for feeding the sheep, and reduced the time spent in rangelands. These changes could be partly explained as a response to decreasing gross margins per livestock unit and the deployment of policies fostering the use of agriculture-based feed resources. Despite these general trends, the variability among pathways of change was wide. Few farms kept using rangelands by focusing on sheep or camel production. Small sheep farms intensified the use of off-farm feeds in the north of the study area, where ecological conditions favoured agriculture. Feed supplementation allowed herd size and animal production to increase, with a substantial risk of susceptibility to market fluctuations. The economic results showed that camel farming combined with small ruminant species can lead to a similar profitability to, or even higher than, large farms that focus solely on sheep and rely on feed supplementation. We conclude that the current situation of livestock farming in arid rangelands remains fragile and their long-term viability is uncertain.

© 2023 The Authors. Published by Elsevier B.V. on behalf of The Animal Consortium. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

### Implications

Feed supplementation has become a common practice in arid regions, but rangelands still constitute an important resource for livestock. Feed supplementation allows herd size and animal productivity to increase but makes farms susceptible to market fluctuations. The optimal balance between the use of rangelands and off-farm feeds depends on local ecological conditions. Camel farming seems a promising strategy to the sustainable use of arid rangelands while maintaining farm profitability.

\* Corresponding author at: Laboratory of Economy and Rural Societies, Arid Regions Institute of Medenine (IRA), Gabes University, 4119 Medenine, Tunisia.

E-mail address: [rjihouda@yahoo.fr](mailto:rjihouda@yahoo.fr) (H. Rjili).

### Introduction

Rangelands cover about 79 million km<sup>2</sup> of the Earth's surface, of which 43% is classified as arid or semi-arid (approx. 34 million km<sup>2</sup>; International Livestock Research Institute-ILRI, 2021). Many pastoral communities across the world directly depend on rangelands for their livelihoods, particularly in arid and semi-arid regions that hold 46% of global livestock production on rangelands (ILRI, 2021). Pastoral communities have historically used rangelands while maintaining an equilibrium between stocking rates and the provision of regulating ecosystem services, such as C and N storage and soil retention and formation, among others (Oñatibia et al., 2015; Fan et al., 2019; Khosravi Mashizi et al., 2019). However, livestock farming systems in these regions have

substantially changed in the last decades due to a variety of global, regional and local drivers.

Similarly to other arid and semi-arid Mediterranean countries, livestock feeding in Tunisia was based on rangeland resources until the 1970s (Abaab and Genin, 2004). Since then, farming systems have undergone major changes such as agricultural mechanisation and the expansion of rainfed cereals, fruit, and olive trees. These changes were driven by policy, market, social and climate factors, fostering livestock-agriculture integration (Ben Salem, 2011; Alary et al., 2019; Rigolot et al., 2019). In Central and Southern Tunisia, these integrated systems have partially replaced the traditional rangeland-based pastoral systems, resulting in transhumant farmers adapting to a sedentary lifestyle (Elloumi et al., 2011; Ammar et al., 2011; Gaddour et al., 2013). Despite these general trends, rangelands still remain an important feed resource for livestock (Bencherif, 2013; Neffati, 2020; Steinfeld et al., 2006) and can be key for the future of livestock farming in the Maghreb (i.e. region of North Africa bordering the Mediterranean Sea, including Algeria, Libya, Mauritania, Morocco, Tunisia, and the disputed territory of Western Sahara). Unlike many other world regions (especially Europe), sheep populations have increased in the last few decades in the Maghreb (Deleule, 2016; Belanche et al., 2021). In the particular case of Tunisia, animal numbers have grown from 8.1 million in 1990 to 8.6 million in 2019, mainly due to sheep (FAOSTAT, 2019). Arid rangelands represent two thirds of national rangelands and provide 20–60% of livestock feed requirements (ILRI, 2021; Jaouad et al., 2022).

General overviews of livestock farming evolution are useful for analysing the main development pathways but fail to account for variability across farms. Exploring the diversity of changes provides evidence of farming heterogeneity, which helps in understanding the range of farmers' reactions to common and specific drivers of change at different scales (e.g., García-Martínez et al., 2009; Ryschawy et al., 2014; Muñoz-Ulecia et al., 2021). Furthermore, becoming aware of past changes can help in designing more effective policies to promote resilient farming in future (Valbuena et al., 2015). However, studies covering temporal dynamics of farming in arid and semi-arid rangelands are scarce, particularly in Maghreb countries (Falconnier et al., 2015; Vall et al., 2017). In Tunisia, several studies have described livestock farm types at specific times (Nefzi, 2012; Jeder et al., 2013; Ibdidhi et al., 2018) but, to our knowledge, no study has analysed the evolution of farming systems, neither the context conditions that may explain evolution patterns (Muñoz-Ulecia et al., 2021). This research bridges this gap by studying changes in livestock farming and geographical patterns of change over a 15-year period.

The objective of this study was twofold: (i) to analyse the main changes that livestock farms in Tunisian arid rangelands have undergone between 2004 and 2019; (ii) to explore potential geographical patterns explaining these changes. We discuss the outcomes of our study in relation to socio-economic and policy factors in the region.

## Material and methods

### Study area

The study was conducted in the El Ouara rangelands in SE Tunisia, which cover 564 000 ha in the Tataouine (Municipalities of Tataouine South, Tataouine North, Rmada and Smar) and Medenine (Benguarden Municipality) governorates (Supplementary Fig. S1). "El Ouara" means "where it is difficult to live", which reflects the harsh climate conditions in our study area. It is a transition zone between Mediterranean woodlands and scrub and xeric scrubland ecological zones (Olson and Dinerstein, 2002) in a Hot Desert Cli-

mate (Köppen climate classification: 175.3 mm annual average rainfall for 2004–2020). The dry season is variable and ranges from 8 to 12 months. Temperatures are mild in winter (January average temperature: 8.4 °C) and very high in summer (August average temperature: 37.2 °C). El Ouara is one of the few common-land rangelands in Tunisia whose main use is still livestock grazing and is, therefore, a relevant example of arid rangelands in Tunisia. The residence place of farmers is generally located in urban areas, 30–120 km away from the rangelands. There are only a few small rural communities near El Ouara, mainly in the Tataouine governorate.

Globally, livestock farming in the study area has changed considerably in recent decades. Traditional pastoral farms mainly dependent on rangeland resources evolved to smallholder mixed livestock-agriculture farms. In these mixed farms, rangeland feed resources are usually complemented with rainfed barley crop residue (stubble, straw, etc.) and agro-industrial by-products (mainly olive pomace) (Ben Salem, 2011; Ibdidhi et al., 2018; Nefzaoui, 2004). Currently, the area accounts for around 19% of the camels, 6% of the goats, and almost 5% of all the sheep in Tunisia (South Development Office, 2018). The region has a negative migratory balance (approximately –6% of the population from 2004 to 2014; official data of the National Institute of Statistics of Tunisia, 2014). Most migrants were young rural people who moved to coastal cities or abroad to seek employment in sectors that offer better labour opportunities than agriculture (Castagnone et Termine, 2021; REACH and Mercy Corps, 2018).

The evolution of employment, land use and livestock heads during the study period differed among municipalities (Supplementary Fig. S2). Although Tataouine governorate experienced a negative net migration, its population grew by 5%. Its main economic activity moved to services and construction, producing a sharp decrease (approx. –70%) in the percentage of people working in agriculture, also observed (–43 to –86% depending on the municipality) in the total cereal crops area (mainly rainfed). However, sheep numbers increased in Tataouine South and North municipalities but lowered in Rmada and in Smar, and the total camel numbers decreased. The areas with tree plantations remained constant (Tataouine South and Rmada) or slightly increased (Tataouine North and Smar). Contrarily to the Tataouine governorate, Benguarden municipality in Medenine Governorate underwent very high population growth (49%). Economic activity moved to construction, while the number of people in the agricultural sector lowered (–59%). Nevertheless, the cereal crop area increased by 66% and tree plantations, which were anecdotal in 2006 (i.e., 1 110 ha), multiplied fifty-fold by 2019. Conversely, the total sheep numbers lowered (30 and 35% respectively), while the total camel numbers slightly rose (1%).

### Farmer survey design and implementation

Data were collected in 2004 and 2019 by a face-to-face survey to farmers. The questionnaire included information about farm structure, management, labour, economic performance, and farmers' socio-economic characteristics. All the farmers using El Ouara rangelands (according to a list drawn up by the Tunisian Union of Agriculture and Fisheries) were surveyed in 2004, totalling 413 farmers (117 for Tataouine South, 90 for Tataouine North, 49 for Rmada, 43 for Smar and 114 for Benguarden). A group of 120 farmers was randomly selected from this sample and contacted again in 2019. Of the 120 farmers, 22 could not be found, 16 had given up livestock farming and nine had passed away without a successor. This left a final sample of 73 farmers who were surveyed both in 2004 and 2019: 41 farms in the Tataouine governorate (13 in Tataouine North, 11 in Tataouine South, 11 in Smar, 6 in Rmada)

and 32 farms in the Medenine governorate (in the Benguarden municipality).

### Data analysis

The analytical methodology is divided into the following four steps:

1. Analysis of the average changes in farms.
2. Analysis of evolution of farms by identifying farm typologies both in 2004 and 2019.
3. Analysis of evolution of farms by identifying the different change pathways followed by farms.
4. Analysis of geographical patterns of typologies and change pathways.

### Analysis of the average changes in livestock farms

We determined the changes in livestock farms during the study period by analysing 22 variables that defined: farm structure, farm management and labour, and farm economic performance (Table 1). Variables were either obtained directly from questionnaires or calculated (i.e., ratios per livestock unit (LU)). The applied LU conversion factors were 1 for camels and 0.15 for both sheep and goats (Agripedia, 2012). All the economic variables were converted into 2019 constant Tunisian dinar (TD, local currency). Differences between years were evaluated by ANOVA, Kruskal-Wallis and chi-square tests depending on the type of variables (i.e., con-

tinuous or categorical) and their distribution (i.e., normal or non-normal). These statistical tests were carried out using the R software (R Core Team, 2019).

### Analysis of the evolution of livestock farms

We analysed the evolution of farms using the analytical method proposed by Doledec and Chessel (1987) and modified by Gibon et al. (1999). This method has been applied to analyse the evolution of farming systems (e.g., García-Martínez et al., 2009; Muñoz-Ulecia et al., 2021), since it allows to analyse differences among farms on each study date, as well as differences in the changes that they had experienced between dates (Gibon et al., 1999). We analysed these two types of changes: intrafarm and interfarm changes (terms used by Gibon et al., 1999). In the intrafarm analysis ("farm typologies" henceforward), farms were compared to the average farm separately on each date. This analysis did not account for the effect of time; it can be considered a farm typology analysis on each date. The interfarm analysis ("change pathways" henceforward) explored differences per farm once the average trend of change (time-dependent) had been eliminated. The method is described in detail below.

### Definition of variables used

We specifically considered 12 continuous variables of the 22 described in Table 1 to be the key descriptors of farm structure and management, including farm structure aspects (i.e., Olive tree area, Cereal area, Herd LU, Camel LU, and Sheep LU/Herd LU), farm

**Table 1**  
Pastoral livestock farm variables considered in the analysis, their nature, description, average values, and SD across years.

Category	Variable ( <sup>1</sup> Cont./Cat.)	Description	2004 Mean ± SD	2019 Mean ± SD
Farm structure	Olive tree area (Cont.)	Farm area used for olive trees (ha).	4.5 <sup>a</sup> ± 6.2	5.8 <sup>b</sup> ± 13.4
	Cereal area (Cont.)	Farm area used for cereal crops (ha).	0.8 <sup>a</sup> ± 2.6	2.3 <sup>b</sup> ± 3.2
	Herd LU (Cont.)	Total livestock units (incl. sheep, goat and camels). Livestock Units (LU) coefficients were 1 for camels and 0.15 for sheep and goats.	41.0 ± 41.5	58.1 ± 67.1
	Camel LU (Cont.)	Number of LU of camels.	17.4 <sup>A</sup> ± 30.4	16.1 <sup>B</sup> ± 36.4
	Sheep LU (Cont.)	Number of LU of sheep.	18.4 ± 19.8	34.7 ± 40.8
	Goat LU (Cont.)	Number of LU of goats.	5.2 ± 5.4	7.3 ± 10.3
	Sheep LU/Herd LU (Cont.)	LU of sheep per total herd LU (%).	54.8 <sup>A</sup> ± 27.6	68.1 <sup>B</sup> ± 29.5
	Tractor ownership (Cat.)	Tractor ownership (yes/no).	23.8%	23.8%
	Wells ownership (Cat.)	Wells ownership (yes/no).	20.5%	21.9%
	Car's ownership (Cat.)	Car's ownership (yes/no).	68.5%	84.9%
	Tank ownership (Cat.)	Tank ownership (yes/no).	23.3%	19.2%
Farm management and labour	WU hired/LU (Cont.)	Non-family work units hired per total livestock units. A work unit (WU) is equivalent to the work of one person, full time, for one year.	0.0 ± 0.0	0.0 ± 0.0
	Rangeland's period (Cont.)	Number of months per year that the herd grazes in rangelands.	10.0 <sup>a</sup> ± 2.5	8.7 <sup>b</sup> ± 3.6
	Transhumance period (Cont.)	Number of months per year that the herd is in transhumance.	3.6 <sup>A</sup> ± 2.6	2.1 <sup>B</sup> ± 1.8
	Second activity	Farmer has other economic activity than farming (yes/no).	28.8%	50.7%
Farm economic performance	Feeding cost/LU (Cont.)	Sum of costs of grains, concentrates, forages and straw use per year per livestock unit (in Tunisian Dinar-TD).	172.7 <sup>A</sup> ± 113.5	250.1 <sup>B</sup> ± 86.5
	Guarding cost/LU (Cont.)	Costs of non-family hired labour for herd guarding per LU.	79.3 <sup>A</sup> ± 102.5	148.4 <sup>B</sup> ± 124.2
	Transport cost/LU (Cont.)	Costs of hired transport to transport feed to the herd when it is in the rangelands and to move animals to rangelands and markets per LU.	24.2 <sup>A</sup> ± 34.7	11.1 <sup>B</sup> ± 28.6
	Water cost/LU (Cont.)	Costs of purchased water and the transport cost (when the well is far from the farm) per LU.	8.4 <sup>a</sup> ± 15.2	6.6 <sup>b</sup> ± 13.0
	Total output (Cont.)	Total income obtained from livestock products in TD.	15 391.7 ± 16 465.4	26 898.7 ± 27 187.7
	GM (Cont.)	Gross margin. It is total output minus livestock costs (feeding, guarding, transport, water and veterinary) in TD.	5 012.7 ± 8 226.7	5 069.5 ± 9 772.8
GM/LU (Cont.)	Gross margin per livestock unit in TD.	119.2 <sup>a</sup> ± 149.8	57.6 <sup>b</sup> ± 153.4	

Abbreviations: LU = Livestock Unit; WU= Work Unit; GM = Gross Margin; TD = Tunisian Dinar (local currency).

<sup>A,B</sup> refer to significant differences ( $P < 0.01$ ) between different change pathways during the study period according to ANOVA, Kruskal-Wallis, or  $\chi^2$  tests depending on the type of variables; <sup>a,b</sup> depict trends ( $P < 0.05$ ).

All the economic figures expressed as 2019 constant Tunisian Dinar (1TD = €0.34, 2019).

<sup>1</sup> Continuous (Cont.) variables are described for each year with average values and SD. Categorical (Cat.) variables are described with the percentage of the sample answering "yes" each year. The same farmers were interviewed in 2004 and 2019 (n = 73).

management and labour aspects (i.e., Work Units (**WU**) hired/LU, Rangeland period, Transhumance period) and input costs (i.e., Feeding cost/LU, Guarding cost/LU, Transport cost/LU, Water cost/LU). Goat LU and Sheep LU were excluded from this analysis because they were highly correlated with Sheep LU/Herd LU, which we considered to be the best proxy to indicate the relative importance of sheep on farms. The three farm economic performance variables: Total output, Gross margin (**GM**), and GM/LU were not considered at this point of the analysis because they do not define farm structure but economic results.

Following Gibon et al. (1999), initially, data were organised in a matrix, in which columns included the  $p$  variables describing farms and which rows included the  $s$  observations (i.e., farms) on the considered  $t$  dates. In our case:  $p = 12$  variables (with normalised values);  $s = 73$  farms;  $t = 2$  dates. Starting from this matrix, the two analyses were carried out as follows.

**Farm typologies in 2004 and 2019.** We built a data table per date where the value of each cell was calculated as  $x_{tsp} - x_{t,p}$  per variable (in columns) representing the deviation of each farm (in rows) to the average of farms per date. A principal component analysis (**PCA**) was performed on both data tables to determine the factors that best explained differences among farms on each date. Then, a K-means cluster analysis (**CA**) was carried out on all the principal components (**PCs**) with eigenvalues above 1 to establish a farm typology. The selection of the number of clusters was based on loss of inertia (in a cluster sum of squares) upon each partitioning of clusters. After establishing the farm typology for 2004 and 2019, we compared the features of the types identified on each date and how individual farms changed, or not, their typologies over the study period.

**Change pathways.** We built a data table with  $p$  columns (i.e., 12 variables with normalised values) and  $s + t$  rows (i.e.,  $s$  observations – 73 farms, on  $t$  dates – 2 time points, 2004 and 2019). The value of each cell was defined as  $x_{tsp} - x_{sp}$ , which represents the deviation to the average of each farm at the two time points. Similarly to the typologies analysis, a PCA was performed. PCs describe the combination of variables that best explained the changes that occurred in farms on the study dates. Then, a CA was performed on the PCs with eigenvalues above 1. Thus, clusters grouped the farms that followed similar change pathways.

Both farm typologies and change pathways were described using the 12 variables considered in the statistical analysis (see above), and 10 complementary variables; two continuous variables (i.e., Sheep LU and Goat LU), five categorical variables (i.e., second economic activity, and tractors, cars, tanks and wells ownership), and three economic performance variables (i.e., Total output, Gross Margin, Gross Margin/LU). The analyses of typologies and change pathways (including the PCA and CA) were done using version 2013 of the XLSTAT software.

#### Analysis of geographical patterns

We analysed the geographical pattern of the evolution of livestock farms by exploring the distribution of farms across municipalities according to their typologies in 2004 and 2019 and change pathways. Differences between municipalities were evaluated by a chi-square test using the R software (R Core Team, 2019).

## Results

### Average changes in livestock farms


Farms changed considerably throughout the study period (Table 1). Cereal area increased by 66% from 2004 to 2019 (Average  $\pm$  SD; 0.8 ha  $\pm$  2.6 ha to 2.3 ha  $\pm$  3.2 ha, respectively;

$P < 0.00$ ), and Olive tree area also increased (4.5 ha  $\pm$  6.2 ha, 5.8 ha  $\pm$  13.4 ha;  $P < 0.03$ ). Sheep LU doubled (18.4 LU  $\pm$  19.8 LU to 34.7 LU  $\pm$  40.8 LU;  $P < 0.07$ ) during the studied period, while Camel LU and Goat LU remained constant, which led to an increase in sheep LU/Herd LU (54.8%  $\pm$  27.6% to 68.1%  $\pm$  29.5%;  $P < 0.00$ ). Work units (**WU**) and hired/LU did not significantly change. We observed a decrease in both the Rangeland grazing period (10.0 months  $\pm$  2.5 months to 8.7 months  $\pm$  3.6 months;  $P < 0.02$ ) and the Transhumance period (3.6 months  $\pm$  2.6 months to 2.1 months  $\pm$  1.8 months;  $P < 0.00$ ). Due to these changes, the total variable costs increased. Feeding cost/LU (172.7 TD  $\pm$  113.5 TD to 250.1 TD  $\pm$  86.5 TD;  $P < 0.00$ ) and Guarding cost/LU (79.3 TD  $\pm$  102.5 TD to 148.4 TD  $\pm$  124.2 TD;  $P < 0.00$ ), which constituted the larger share of the variable costs, also increased. Transport cost/LU more than halved (24.2 TD  $\pm$  34.7 TD to 11.1 TD  $\pm$  28.6 TD;  $P < 0.00$ ), and Water costs slightly decreased (8.4 TD  $\pm$  15.2 TD to 8.0 TD  $\pm$  15.6 TD;  $P < 0.02$ ). In parallel to the rise in variable costs, Total output almost doubled (15391.7 TD  $\pm$  16465.4 TD to 26898.7 TD  $\pm$  27187.7 TD;  $P < 0.06$ ). However, Gross margin (**GM**) remained stable during the study period (5 012.7 TD  $\pm$  8 226.7 TD to 5 069.5 TD  $\pm$  9 772.8 TD;  $P < 0.49$ ) because the GM/LU halved (119.2 TD  $\pm$  149.8 TD to 57.6 TD  $\pm$  153.4 TD;  $P < 0.02$ ).

### Farm typologies in 2004 and 2019

PCA results are provided in the Appendix (Supplementary Table S1). CA resulted at three clusters in both 2004 and 2019 (i.e., “2004 typologies” and “2019 typologies” hereafter). Similar typologies were identified in both dates, although their average features and relative importance evolved over time: (a) camel-focused farms with large tree areas (“Camel farms”, hereafter); (b) large sheep-focused farms with considerable rangelands use (“Large sheep farms”, hereafter); (c) small sheep-focused farms (“Small sheep farms”, hereafter). Typologies are described below according to the 12 variables considered in the analysis of changes (Fig. 1) and 10 other complementary variables (Table 2).

- Camel farms.** In 2004, these farms represented 24.7% of the sampled population and their relative importance decreased during the study period; in 2019, they represented 15.0% of the farms. They were characterised by having the highest Camel LU and Olive tree area, and the lowest Sheep LU/Herd LU of the three typologies. They had the longest transhumance period and the lowest feeding and water costs per LU. From 2004 to 2019, these farms had an increased Camel LU ( $P < 0.01$ ). GM/LU more than halved during the study period ( $P < 0.06$ ), but GM remained constant.
- Large sheep farms.** In 2004, these farms represented 23.3% of the sampled population and their relative importance almost doubled during the study period; in 2019, they represented 49.3% of the farms. They had the highest Sheep and Goat LU and, contrarily to the Small sheep farms (see below), they usually recruited external labour and practised transhumance for longer periods than other typologies. Besides these aspects, they were similar to Small sheep farms in farming system management terms. During the study period, the Cereal area per farm greatly increased ( $P < 0.0001$ ). The Rangeland period remained stable, but the Transhumance period significantly reduced ( $P < 0.00$ ). Both GM and GM/LU remained constant.
- Small sheep farms.** In 2004, these farms represented more than half (52%) the sampled population and their relative importance decreased during the study period; in 2019, they represented 35.6% of the farms. They were characterised for having the lowest Herd LU, Transhumance period, WU hired/LU and Guarding cost/LU of all typologies. Cereal area tripled



**Fig. 1.** Changes observed in the variables used to define pastoral livestock farm typologies across years. Boxplots represent the median (solid horizontal lines), the first and third quartiles (contained in boxes) and outliers (black points). Different letters refer to significant differences between farm typologies per year according to ANOVA. Abbreviations; LU = Livestock Unit WU = work unit TD = Tunisian Dinar (local currency).

**Table 2**  
Description of pastoral livestock farm typologies, average values, and SD across farm typologies for each year.

Category	Variable	2004			2019		
		Camel farms (n = 18)	Large sheep farms (n = 17)	Small sheep- farms (n = 38)	Camel farms (n = 11)	Large sheep farms (n = 36)	Small sheep farms (n = 26)
<b>Farm structure</b>							
	Sheep LU (LU)	21.1 <sup>AB</sup> ± 34.8	33.5 <sup>A</sup> ± 25.1	10.4 <sup>B</sup> ± 9.4	55.3 <sup>AB</sup> ± 67.0	48.5 <sup>A</sup> ± 5.8	6.8 <sup>B</sup> ± 22.0
	Goat LU (LU)	3.7 <sup>A</sup> ± 12.2	9.8 <sup>B</sup> ± 4.4	3.8 <sup>A</sup> ± 3.9	8.1 <sup>AB</sup> ± 9.9	11.0 <sup>A</sup> ± 2.8	2.0 <sup>B</sup> ± 6.7
	Tractor ownership <sup>1</sup> (% of farmers)	27.8	23.5	21.0	54.6 <sup>a</sup>	22.2 <sup>ab</sup>	11.5 <sup>b</sup>
	Cars ownership <sup>1</sup> (% of farmers)	88.9 <sup>a</sup>	76.5 <sup>ab</sup>	55.3 <sup>b</sup>	100.0	83.3	80.8
	Tank ownership <sup>1</sup> (% of farmers)	29.4	33.3	15.8	45.5	16.7	11.5
	Wells ownership <sup>1</sup> (% of farmers)	17.7	22.2	21.1	27.3	16.7	26.9
<b>Farm management and labour (% of farmers)</b>							
	Second Activity <sup>1</sup>	11.8	38.9	34.2	63.6	47.2	57.7
<b>Farm economic performance (TD)</b>							
	Total output	24 812.1 <sup>A</sup> ± 26 371.7	22 705.2 <sup>A</sup> ± 20 107.6	7 657.6 <sup>B</sup> ± 21 478.6	5 0182.7 <sup>A</sup> ± 29 438.7	36 095.3 <sup>A</sup> ± 23 668.3	4 314.0 <sup>B</sup> ± 7 531.0
	GM	7 905.5 ± 10 085.0	6 571.5 ± 9 681.4	2 945.0 ± 8 008.2	8 410.0 <sup>AB</sup> ± 11 400.4	7 101.4 <sup>A</sup> ± 10 354.1	842.7 <sup>B</sup> ± 3 749.7
	GM/LU	99.1 ± 104.7	94.8 ± 177.7	139.0 ± 164.6	36.4 ± 118.3	69.5 ± 161.6	50.2 ± 154.8

Abbreviations: LU = Livestock Unit; GM = gross margin; TD = Tunisian Dinar (local currency).

<sup>A,B</sup> refer to significant differences ( $P < 0.01$ ) between different change pathways during the study period according to ANOVA, Kruskal-Wallis, or  $\chi^2$  tests depending on the type of variables; <sup>a,b</sup> depict trends ( $P < 0.05$ ).

All the economic figures expressed as 2019 constant Tunisian Dinar (1TD = €0.34, 2019).

<sup>1</sup> Categorical variables are described with the percentage of the sample answering “yes” each year.

from 2004 to 2019 ( $P < 0.00$ ) and was the highest of all the typologies in 2019. The Herd size of this farm type almost halved throughout the study period and the number of camels in herds lowered ( $P < 0.05$ ). The Rangeland period also halved ( $P < 0.00$ ) with the consequent increase in Feeding cost/LU (18.7%;  $P < 0.05$ ) and a cut in Transport cost/LU ( $P < 0.00$ ). As in the other two farming system typologies, GM/LU decreased during the study period ( $P < 0.04$ ) but, conversely, GM also reduced ( $P < 0.02$ ).

The fact that similar farm typologies were found in 2004 and 2019 does not mean that individual farms continued with the same typology during that period. Fig. 2 shows how individual farms shifted among the typologies between 2004 and 2019. Around half ( $n = 35$ ) of the farms fell in the same typology in both 2004 and 2019, which was more likely to happen in the Large sheep farms than in the other two typologies. Most of the Large sheep farms that did not remain in the same category shifted to the Small sheep farms (24%). On the contrary, half of the Small sheep farms in 2004 became Large sheep farms in 2019, and only a very low percentage (5%) became Camel farms. Finally, a similar proportion (approx. 30%) of the 2004 Camel farms shifted to both the Small sheep farms and Large sheep farms.

*Farms' change pathways*

The PCA resulted in five PCs with an eigenvalue above 1 that explained 69% of total variance and represented the major trends of change in the sampled farms (Supplementary Table S2). Based on these five PCs, the CA resulted in four farms' change pathways, which are described according to the 12 variables used in the analysis of pathways (Fig. 3) and the other 10 complementary variables (Table 3).

*Sheep intensification pathway*

This pathway was followed by 20.5% of the sampled population. These farms were characterised by the largest increase in Sheep LU/Herd LU and Feeding cost/LU of all the pathways, and by a sharp

reduction in the Rangeland period. This reduction in herd movement led both Transport and Water costs/LU to lower. All these changes brought about an increase in Total output, although GM and GM/LU decreased. They also increased the agricultural area, particularly the Olive tree area. Finally, these farms showed the highest decrease in household size (−50%) and the highest increase in practicing a second economic activity (80% of the farms started a second activity besides farming).

*Non-sheep extensification pathway*

This pathway was followed by 20.5% of the sampled population and was the only one to show an extension in both rangelands and transhumance periods. In parallel, feeding costs lowered by an average of 37.6%, whereas Guarding cost/LU increased. Farms in this pathway showed the highest increase in herd size of all the pathways, which was directly related to an increase in both Camel LU and Goat LU which, in turn, led to a decrease in Sheep LU/Herd LU. Average farm output and GM increased despite GM/LU lowering. Olive tree area remained more or less constant but, like the other pathways, Cereal area increased. Finally, more than half the farms (66.7%) started secondary activities during the study period.

*Sheep rangeland pathway*

It was followed by 26.1% of the sampled population. These farms restructured their herd by reducing the number of camels and slightly increasing the number of small ruminants (particularly sheep), while keeping the total herd size stable. In parallel, the Rangeland period slightly increased, but the Transhumance period sharply decreased. Workforce hired and, consequently, the Guarding cost/LU, increased. Total output increased slightly, but GM and GM/LU went down. Cereal area increased, but Olive tree area reduced (3.7 ha in 2019). Almost 40% of farms started a secondary activity.

*Stable herd structure pathway*

This pathway was followed by 32.9% of the sampled population. It grouped the farms that remained relatively stable in terms of herd size and composition. However, unlike the other pathways,





Fig. 2. Shifting among pastoral livestock farm typologies between 2004 and 2019. The width of the grey linking lines is proportional to the number of farms.



**Fig. 3.** Trends of change observed in the variables defining pastoral livestock farm change pathways during the study period. Boxplots represent the median (solid horizontal lines), the first and third quartiles (contained in boxes) and outliers (black points) of the distribution of the variables. Different letters refer to significant differences between pathways for each variable according to ANOVA. Abbreviations: LU = Livestock Unit; WU = work unit; TD = Tunisian Dinar (local currency).

**Table 3**

Trends of changes observed in pastoral livestock farm pathways. Average values and SD of the value of each variable in 2019, minus the average of this variable in the two years across pathways.

Category	Variable	Sheep intensification pathway (n = 15) Mean ± SD	Non-sheep extensification pathway (n = 15) Mean ± SD	Sheep rangelands pathway (n = 19) Mean ± SD	Stable herd structure pathway (n = 24) Mean ± SD
<b>Farm structure</b>					
	Sheep LU (LU)	10.7 <sup>ab</sup> ± 16.9	20.2 <sup>a</sup> ± 27.0	5.4 <sup>ab</sup> ± 14.8	1.2 <sup>b</sup> ± 15.7
	Goat LU (LU)	0.0 <sup>a</sup> ± 2.7	4.3 <sup>B</sup> ± 6.1	2.1 <sup>AB</sup> ± 7.8	-1.1 <sup>A</sup> ± 2.8
	Tractor ownership <sup>1</sup> (% of farmers)	33.3	33.3	47.4	29.2
	Cars ownership <sup>1</sup> (% of farmers)	40.0	46.7	36.8	33.3
	Tank ownership <sup>1</sup> (% of farmers)	40.0	33.3	26.3	33.3
	Wells ownership <sup>1</sup> (% of farmers)	46.7	26.7	31.6	25.0
<b>Farm management and labour (% of farmers)</b>					
	Second Activity <sup>1</sup>	80.0	66.7	36.8	25.0
<b>Farm economic performance (TD)</b>					
	Total output	6 468.4 ± 12 534.3	14 613.2 ± 16 875.5	2 463.0 ± 10 571.1	2 374.3 ± 9 650.6
	GM	-604.1 <sup>ab</sup> ± 3 709.7	2 694.8 <sup>a</sup> ± 7 059.1	-3 001.3 <sup>b</sup> ± 4 410.6	1 155.7 <sup>a</sup> ± 5 278.9
	GM/LU	-36.9 <sup>ab</sup> ± 101.0	-70.9 <sup>a</sup> ± 120.6	-59.2 <sup>A</sup> ± 63.6	20.7 <sup>B</sup> ± 90.6

Abbreviations: LU = Livestock Unit; GM = gross margin; TD = Tunisian Dinar (local currency).


<sup>A,B</sup> refer to significant differences ( $P < 0.01$ ) between different change pathways during the study period according to ANOVA, Kruskal-Wallis, or  $\chi^2$  tests depending on the type of variables; <sup>ab</sup> depict trends ( $P < 0.05$ ).

All the economic figures are expressed in 2019 constant Tunisian Dinar (1TD = €0.34, 2019).

<sup>1</sup> Categorical variables are expressed as the percentage of farms that changed.

Olive tree area showed a major decrease. Herd management changed to a certain extent with a reduction in the Rangeland and Transhumant periods and, consequently, Feeding cost/LU increased and

Transport costs/LU lowered. Both Total output and GM increased, and, unlike the other pathways, GM/LU also increased. Finally, the farms had the lowest increase in secondary activities (25%).



**Fig. 4.** Geographical distribution of pastoral livestock farms according to their typologies in 2004 and 2019 and their change pathways. For each municipality, pie charts represent the proportion of farms in each typology across years and in each pathway.

**Geographical patterns**

Our results showed statistical differences across municipalities in the distribution of farms typologies both in 2004 (Chi-square test  $P$ -value  $<0.03$ ) and 2019 (Chi-square test  $P$ -value  $<0.01$ ). In 2004, most Camel farms were located in Benguarden municipality. In the other municipalities, Small sheep farms represented around three quarters of farms, with the remaining quarter being mostly Large Sheep farms (Fig. 4). This picture had changed in 2019. While most Camel farms were still located in Benguarden, Small Sheep farms tended to be located in the Northern part of the study area, and Large Sheep farms in the Southern part. We found no statistical differences in the distribution of farm change pathways across municipalities (Chi-square test  $P$ -value = 0.28). However, we observed a consistent trend within municipalities where farms following the Sheep intensification pathway were located in the Northern part of each municipality. Note that no Sheep intensification farms were located in Rmada municipality.

**Discussion**

This study allowed us to explore the general evolution of livestock farms over a 15-year period in a representative arid rangeland of South Tunisia. We also considered the diversity of farm typologies and change pathways, being one of the few studies ana-

lysing livestock farming dynamics in arid rangelands (Falconnier et al., 2015). Below we discuss in detail the farm typologies and change pathways, the geographical patterns, and how they might be related to socio-economic factors.

*General change: increase in herd size and cereal area, sheep specialisation, and reduction of time spent in rangelands*

Most farms evolved in the same direction for four aspects: increase in herd size, increase in cereal area, stronger orientation to sheep production, and (limited) reduction in transhumance and time spent in rangelands. The increase in herd size aligns with a general trend observed in other arid and semi-arid regions of Tunisia, North Africa and elsewhere (Jemaa, 2016; Maatougui, 2000; Mohamed et al., 2021). This is likely a response to the reduction of GM/LU due to the low price of lambs (Bencherif, 2013), and the availability of supplementary feed resources, as discussed below. The increase in herd size was due to sheep numbers while goat numbers decreased. These findings confirm a trend that started one decade before the study was conducted (Ben saad and Bourbouze, 2010). Goats are of lower commercial interest and are mostly kept for milk home-consumption.

In most cases, the increase of herd size was accompanied by supplementary on-farm feeds (i.e., rainfed agriculture by-products, mainly thatch and straw) or purchased, which increased



the variable costs per LU and, in turn, further reduced GM/LU. Feed intensification is a common trend observed across farming systems, livestock species and world regions (Powell et al., 2004; Vall et al., 2017; Godde et al., 2018). However, El Ouara farms used less supplementary feeds compared to other regions in Tunisia (e.g., centre of the country: Ibdhi and Ben Salem (2018); Jemaa (2016) and Maghreb areas (Bensmira, 2017). In line with Abdelguerfi and El Hassani (2011) and Hadbaoui et al. (2020), we found a large integration of agriculture and livestock activities which was almost inexistent some decades ago (Abbas, 2014; Nasr, 2004) and very limited at the beginning of the study. Higher use of cereals and agriculture by-products for livestock feeding has improved animal performance but also increased cereal price volatility at national and international levels, and therefore production costs. In Tunisia (Elloumi, 2015) and other developing countries, the dependence on agricultural commodities produced elsewhere has been pointed out as one of the causes of economic instability and social crisis in the recent past (Behnassi and El Haiba, 2022; Jayasuriya et al., 2012; Mittal, 2009), which are likely to be exacerbated in the context of climate change (Vesco et al., 2021).

The average time that herds spend in rangelands was reduced. Previous studies have shown that the contribution of rangelands to small ruminant feeding has decreased in Tunisia in the last three decades (Ibdhi and Ben Salem, 2018). These studies point to overgrazing and rainfall reduction as key drivers for the reduction of the use of rangelands. We can neither confirm nor deny these drivers. Yet, when analysing the different change pathways, our results revealed that some of farms maintained the use of rangelands. In addition, our results showed a general increase in farm machinery and infrastructures (i.e., tractors, cars, wells). Tractors and cars increase might be related to crop production, but they can also be used to transport water and feed, allowing livestock to remain on rangelands during dry periods (Nefzaoui et al., 2012).

In most typologies, guarding costs increased due to rising shepherd salaries (Selmi et al., 2018). According to informal conversations with farmers, this is a consequence of the lack of specialised shepherds. This is probably due to emigration that particularly involves young people (Richard, 2006). Livestock management in harsh environments, such as El Ouara rangelands, requires specialised shepherds with in-depth knowledge of resource distribution across space and time (Bourbouze, 2018; Selmi and Elloumi, 2007). Lack of specialised shepherds is also a common problem in many other world regions and is generally considered one of the main challenges that future pasture-based systems will face (Morales et al., 2019; Paniagua, 2019).

Finally, the results showed that many farmers (between 25 and 47 out of 120) quit their activity during the study period, which is in line with the decrease in agricultural employment in Tunisia (National Institute of Statistics of Tunisia, 2018). During informal conversations, several farmers pointed out that lack of family labour and/or economic resources to make farm investments are the main reasons for farming abandonment. It is not known to what extent this process will continue in future, but the current scenario of stagnated profits and young people migration will not help halting the abandonment of farming.

#### *Farm typologies and change pathways*

Regardless of the pathway followed by farms, feeding costs per LU increased mainly due to purchased supplementary feed. In two pathways, this increase was accompanied by a reduction in the use of rangeland (regardless of the evolution of the number of sheep). The economic performance analysis showed that the strategy which focused on sheep maintaining rangeland use led to a sharp drop in GM, which was not observed in other pathways. This result

should be confirmed with more accurate farm economic assessments, including other factors affecting household economic performance such as labour and access to land. However, if it holds true, it does not bode well for the future of rangeland-based sheep farming in El Ouara.

On the contrary, the non-sheep extensification pathway, which focused on camels with a limited number of sheep and maintained transhumance practices in rangelands, was the most profitable pathway. This result suggests that mixed camel-sheep farming could obtain similar or higher profitability than large sheep farms relying on feed supplementation. Studies in other African drylands have found this combination profitable (Behnke, 2021; Faye and Bonnet, 2012; Ratemo et al., 2020). Although the Tunisian government has set up a national programme for camel research and development, production remains a traditional activity that relies on arid rangelands with low reproductive efficiency (Moslah et al., 2004; Jemli et al., 2018). Hence, according to some experts (Faye et al., 2014), future camel farming will depend on the sector's ability to improve herd productivity and market channels, which still are largely informal.

#### *Geographical patterns*

Our study shows a clear relation between farms' geographical location and pathways of change, which became stronger in 2019 and showed a North-South gradient. This pattern might be explained by several interrelated socio-economic and ecological factors. On the one hand, the fact that small sheep farms (in 2019) and the intensification pathway were located mainly in the north might respond to differential ecological features of El Ouara, i.e., "Jeffara" plains in the north far more favourable for agriculture (Guillaume, 2009). Therefore, in favourable agricultural areas, small sheep farms may intensify by utilising agriculture-based feeds, supported by specific sectoral policies (Elloumi, 2006). The harder ecological conditions in the south do not allow the development of crop farming, which explain why farmers increase herd size and rangeland use. On the other hand, camel farms were almost exclusively located in Benguarden municipality in both 2004 and 2019. Benguarden holds the largest camel population of all the municipalities in Tunisia (ODS, 2018). This is linked with socio-cultural factors: the "Twazin" tribe, mainly located in the Medenine governorate (Benguarden municipality), has historically raised camels (Moslam and Megdiche, 1989). In addition, camel farming is supported by higher consumption and demand of camel meat in southern regions (Moslah et al., 2004; Trabelsi, 2016).

#### *Limitations*

Survey-based results have limitations that should be considered. Collecting accurate on-farm data from face-to-face questionnaires applied to smallholder farmers in developing countries is challenging. Farmers do not record any animal or farm performance data and, therefore, aspects related to animal productivity, feed intake or feed production could not be analysed. Hence, we should acknowledge that the economic data are based on farmers' estimations, not on accurate accounting. In addition, few official data are available on the socio-economic and environmental factors at the municipality level (where our study focuses on). Therefore, we cannot establish the causal relations between socio-economic and environmental drivers, as other studies do (e.g. Brown et al., 2019; Caron and Hubert, 2000; Muñoz-Ulecia et al., 2021). For these reasons, we decided to limit our discussion to general aspects, for which we point out plausible explanations for the observed changes.

The above-mentioned limitations are common in developing countries and might be one of the reasons why studies of temporal dynamics are scarce in many world regions. This does not limit the interest of our study, which offers relevant information about the changes taking place in poorly known farming systems.

#### *Implications for the future of livestock farms in arid rangelands*

Two main implications for the future of livestock farms in arid rangelands derive from our study. Firstly, feed supplementation has become widespread, and additional feed resources (produced on or off farm) will very likely drive the future development of livestock farms in El Ouara and other arid rangelands elsewhere. Feed supplementation might allow herd size, animal productivity and farm profitability to increase, which is particularly key in a climate change context, where reduced primary production in rangelands is expected (Godde et al., 2019). However, this very much depends on feed production costs and market prices. Intensification of feed management makes farms more susceptible to external markets and commodity prices, which will be a key issue for farm sustainability and resilience in developing countries (e.g., Lorent et al., 2009; Cortner et al., 2019) and elsewhere. Moreover, increasing herd sizes may bring about further rangeland overexploitation, which has been observed in different areas of the Maghreb (Bechchari et al., 2014; Bencherif, 2018). The optimum balance between sustainable use of rangelands and of additional feed resources will also depend on herd structure, management, and location, according to ecological conditions. Field studies are needed to determine the stocking rate that can allow natural rangeland regeneration.

Secondly, the diversification of livestock species appears to be a promising strategy. Specifically, adaptation capacity to harsh environments can make camels crucial in the climate change context of arid regions. However, the introduction of camels on small ruminant farms poses many challenges because it requires expensive upfront investments and knowledge (Volpato and King, 2019). The camel sector is still weak and disorganised, and the valorisation of camel products (milk, meat) is still below its potential, despite some improvements observed in recent years (Faye, 2013; Jemli et al., 2018).

Finally, even if farms have evolved during the study period in response to changing socio-economic conditions, their current situation (i.e., old farmers, decreasing household size and declining farm profits) remains fragile and their long-term viability is uncertain.

#### **Supplementary material**

Supplementary material to this article can be found online at <https://doi.org/10.1016/j.animal.2023.100748>.

#### **Ethics approval**

Not applicable.

#### **Data and model availability statement**

None of the data were deposited in an official repository. The data that support the findings of this study are available upon reasonable request to the corresponding author.

#### **Author ORCIDs**

**H.Rjili:** <https://orcid.org/0000-0001-9136-2813>.

**E. Muñoz-Ulecia:** <https://orcid.org/0000-0002-7153-7660>.

**A. Bernués:** <https://orcid.org/0000-0002-3237-9751>.

**M. Jaouad:** <https://orcid.org/0000-0001-5202-6024>.

**D. Martin-Collado:** <https://orcid.org/0000-0002-2087-961X>.

#### **Author contributions**

**H. Rjili:** Conceptualisation, Investigation, Data curation, Formal analysis, Methodology, Software, Writing – original draft. **E. Muñoz-Ulecia:** Data presentation, Software, Methodology, Writing – review & editing. **A. Bernués Jal:** Writing – review & editing. **M. Jaouad:** Resources, Conceptualisation, Writing – review & editing. **D. Martin-Collado:** Conceptualisation, Methodology, Software, Supervision, Writing – original draft, Writing – review & editing, Validation.

#### **Declaration of interests**

None.

#### **Acknowledgements**

The authors are thankful to the local agricultural authorities (Regional Commissariat for Agricultural Development of both Medenine and Tataouine governorates) for their support in conducting the survey of farmers. A special mention is reserved for farmers who have agreed to answer the questions of the survey.

#### **Financial support statement**

The research leading to these results has received funding from an exchange scholarship of the Ministry of Higher Education and Scientific Research, Carthage University, Tunisia. H. Rjili is inscribed in a PhD programme supported by the Laboratory of Economy and Rural Society of the Institute of Arid Regions, Medenine, Tunisia.

#### **References**

- Abaab, A., Genin, D., 2004. Politiques de développement agropastoral au Maghreb. In *Environnement et sociétés rurales en mutation. Approches alternatives* (ed. Picouet, M., Sghaier, M., Genin, D., Abaab, A., Guillaume, H., Elloumi, M.). IRD éditions, Paris, France, pp. 341–358.
- Abbas, K., 2014. Enjeux du développement de l'élevage ovin en zones céréalières semi-arides algériennes. Papier présenté au 8ème séminaire du sous-réseau FAO-CIHEAM sur les systèmes de production, 11–13 June 2013, Tanger, Morocco, pp. 469–475.
- Abdelguerfi, A., El Hassani, T.A., 2011. Interactions entre les systèmes de culture céréalière et les zones pastorales comme base pour le développement durable de l'agriculture dans les pays méditerranéens. In: Lemaire, G., Hodgson, J., Chabbi, A. (Eds.), *Grassland Productivity and Ecosystem Services*. CABI Digital Library, Wallingford, UK, pp. 261–271.
- Agripedia, 2012. Coefficients de conversion en unités de gros bétail, Retrieved on 7 October 2020 from <https://agripedia.ch/terminologie/fr/annexes/annexe-5-coefficients-de-conversion-en-unites-de-gros-betail/>.
- Alary, V., Moulin, C.H., Lasseur, J., Aboul-Naga, A., Sraïri, M.T., 2019. The dynamic of crop-livestock systems in the Mediterranean and future prospective at local level: A comparative analysis for South and North Mediterranean systems. *Livestock Science* 224, 40–49. <https://doi.org/10.1016/j.livsci.2019.03.017>.
- Ammar, H., Bodas, R., Younes, M.B., López, S., 2011. Goat breeding systems in the South of Tunisia (Tataouine). Economic, social and environmental sustainability in sheep and goat production systems. Zaragoza CIHEAM/FAO/CITA/DGA. *Options Méditerranéennes: Série A. Séminaires Méditerranéens* 100, 283–288.
- Bechchari, A., El Aich, A., Mahyou, H., Baghdad, M., Bendaou, M., 2014. Analyse de l'évolution du système pastoral du Maroc oriental. *Revue d'élevage et de médecine vétérinaire des pays tropicaux* 67, 151–162.
- Behnassi, M., El Haiba, M., 2022. Implications of the Russia-Ukraine war for global food security. *Nature Human Behaviour* 6, 754–755. <https://doi.org/10.1038/s41562-022-01391-x>.
- Behnke, R.H., 2021. Grazing Into the Anthropocene or Back to the Future? *Frontiers in Sustainable Food Systems* 5, 116.
- Belanche, A., Martín-Collado, D., Rose, G., Yáñez-Ruiz, D.R., 2021. A multi-stakeholder participatory study identifies the priorities for the sustainability of the small ruminants farming sector in Europe. *Animal* 15, 100–131. <https://doi.org/10.1016/j.animal.2020.100131>.

- Ben saad, A., Bourbouze, A., 2010. Options Méditerranéennes en ligne – Collection numérique – Les nouveaux visages du pastoralisme moderne du grand sud tunisien. Retrieved on 10 February 2022 from <https://om.ciheam.org/article.php?IDPDF=801270>.
- Ben Salem, H.B., 2011. Mutations des systèmes alimentaires des ovins en Tunisie et place des ressources alternatives 12. In Mutations des systèmes d'élevage des ovins et perspectives de leur durabilité (ed. Khlij, E., Ben Hamouda, M., Gabiñ a D.). Options Méditerranéennes: Série A. Séminaires Méditerranéens 97, 29–39.
- Bencherif, S., 2013. L'élevage agropastoral de la steppe algérienne dans la tourmente: enquêtes et perspectives de développement. Mondes en développement 161, 93–106. <https://doi.org/10.3917/med.161.0093>.
- Bencherif, S., 2018. Origines et transformations récentes de l'élevage pastoral de la steppe algérienne. Revue internationale des études du développement 236, 55–79.
- Bensmira, Z., 2017. Contribution à l'étude des systèmes d'élevages ovins et gestion des parcours steppiques de Ras El Ma de Sidi Bel Abbes (Algérie occidentale). PhD Thesis, Université Djillali Liabes, Sidi Bel Abbes, Algérie.
- Bourbouze, A., 2018. Les grandes transformations du pastoralisme méditerranéen et l'émergence de nouveaux modes de production. In Animal health and livestock, Mediterranean perspectives. Watch Letter - Lettre de veille du CIHEAM 39, 7–12.
- Brown, P., Daigneault, A., Dawson, J., 2019. Age, values, farming objectives, past management decisions, and future intentions in New Zealand agriculture. Journal of Environmental Management 231, 110–120. <https://doi.org/10.1016/j.jenvman.2018.10.018>.
- Caron, P., Hubert, B., 2000. De l'analyse des pratiques à la construction d'un modèle d'évolution des systèmes d'élevage: application à la région Nordeste du Brésil. Revue d'élevage et de médecine vétérinaire des pays tropicaux 53, 37–53. <https://doi.org/10.19182/remvt.9763>.
- Castagnone, E., Termine, P., 2021. Migration des jeunes ruraux Méditerranéens: déterminants socio-économiques, défis et opportunités pour l'élaboration de politiques ciblées. In MediTERRA: Migrations et développement rural inclusif en Méditerranée (ed. Presses de Sciences Po). La Librairie Européenne, Bruxelles, Belgique, pp. 147–167.
- Cortner, O., Garrett, R.D., Valentim, J.F., Ferreira, J., Niles, M.T., Reis, J., Gil, J., 2019. Perceptions of integrated crop-livestock systems for sustainable intensification in the Brazilian Amazon. Land Use Policy 82, 841–853. <https://doi.org/10.1016/j.landusepol.2019.01.006>.
- Deleule, M., 2016. Évolution des systèmes d'élevage dans les steppes du Maghreb: enjeux et perspectives. Mémoire Master 2, University of Montpellier, Montpellier, France.
- Doledéc, S., Chessel, D., 1987. Seasonal successions and spatial variables in freshwater environments. I. Description of a complete two-way layout by projection of variables. Acta Oecologica/Oecologia Generalis 8, 403–426.
- Eloumi, M., 2006. Les politiques de développement rural en Tunisie: Acquis et perspectives. Papier présentée au Séminaire sur les Politiques de Développement Rural Durable en Méditerranée dans le Cadre de la Politique de Voisinage de l'Union Européenne, 08–09 février 2006, Le Caire, Egypt. Options Méditerranéennes 71, 55–65.
- Eloumi, M., Selmi, S., Zaibet, L., 2011. Importance économique et mutation des systèmes de production ovins en Tunisie. In Mutations des systèmes d'élevage des ovins et perspectives de leur durabilité (ed. Khlij, E., Ben Hamouda, M., Gabiñ a D.). Options Méditerranéennes: Série A. Séminaires Méditerranéens 97, 11–21.
- Eloumi, M., 2015. Capacité de résilience de l'agriculture familiale tunisienne et politique agricole post révolution. In Accaparement, action publique, stratégies individuelles et ressources naturelles: regards croisés sur la course aux terres et à l'eau en contextes méditerranéens (ed. Centre International De Hautes Etudes Agronomiques). Options Méditerranéennes, CIHEAM, Montpellier, France, pp. 351–366.
- Falconnier, G.N., Descheemaeker, K., Van Mourik, T.A., Sanogo, O.M., Giller, K.E., 2015. Understanding farm trajectories and development pathways: Two decades of change in southern Mali. Agriculture System 139, 210–222. <https://doi.org/10.1016/j.agsy.2015.07.005>.
- Fan, F., Liang, C., Tang, Y., Harker-Schuch, I., Porter, J.R., 2019. Effects and relationships of grazing intensity on multiple ecosystem services in the Inner Mongolian steppe. Science of The Total Environment 675, 642–650. <https://doi.org/10.1016/j.scitotenv.2019.04.279>.
- FAOSTAT, 2019. Crop and livestock products. Retrieved on 10 February 2022 from <http://www.fao.org/faostat/fr/#data/QA>.
- Faye, B., 2013. Camel farming sustainability: the challenges of the camel farming system in the XXIth century. Journal of Sustainable Development 6, 74–82.
- Faye, B., Bonnet, P., 2012. Camel Sciences and Economy in the World: Current Situation and Perspectives. Paper presented at the 3rd conference of the international society of camild research and development, 29 January – 1 February 2012, Muscat, Sultanate of Oman.
- Faye, B., Madani, H., El-Rouili, S.A., 2014. Camel milk value chain in Northern Saudi Arabia. Emirates Journal of Food and Agriculture 26, 359–365.
- Gaddour, A., Najari, S., Abdennebi, M., 2013. Spécificité et diversité des systèmes de reproduction caprine et ovine dans les régions arides Tunisiennes. In Technology creation and transfer in small ruminants: roles of research, development services and farmer associations (ed. Chentouf M., López-Francos A., Ben goumi M., Gabiñ a D.). Options Méditerranéennes: Série A. Séminaires Méditerranéens 108, 477–480.
- García-Martínez, A., Olaizola, A., Bernués, A., 2009. Trajectories of evolution and drivers of change in European mountain cattle farming systems. Animal 3, 152–165. <https://doi.org/10.1017/S1751731108003297>.
- Gibon, A., Balent, G., Olaizola, A., Di Pietro, F., 1999. Approche des variations communales des dynamiques rurales au moyen d'une typologie: cas du versant nord des Pyrénées centrales. Options Méditerranéennes: Série B Etudes de Recherche 27, 15–34.
- Godde, C., Dizyee, K., Ash, A., Thornton, P., Sloat, L., Roura, E., Henderson, H., Herrero, M., 2019. Climate change and variability impacts on grazing herds: Insights from a system dynamics approach for semi-arid Australian rangelands. Global Change Biology 25, 3091–3109.
- Godde, C.M., Garnett, T., Thornton, P.K., Ash, A.J., Herrero, M., 2018. Grazing systems expansion and intensification: drivers, dynamics, and trade-offs. Global Food Security 16, 93–105. <https://doi.org/10.1016/j.gfs.2017.11.003>.
- Guillaume, H., 2009. Mutations agro-pastorales, ruralité et développement dans le Sud-Est tunisien. In: Bonte, P., Eloumi, M., Guillaume, H., Mahdi, M. (Eds.), Développement rural, Environnement et Enjeux territoriaux. Regards croisés Oriental marocain et Sud-Est tunisien. Cérès, Tunis, Tunisie, pp. 19–43.
- Hadbaoui, I., Senoussi, H., Huguenin, J., 2020. Les modalités d'alimentation des troupeaux ovins en steppe algérienne, région de M'Sila: pratiques et tendances. Cahiers Agricultures 29, 28. <https://doi.org/10.1051/cagri/2020027>.
- Ibidhi, R., Ben Salem, H., 2018. Analysis of small ruminants-based farming systems and their interaction with freshwater resources in Tunisia: Interventions for improving water productivity. Annales de l'INRAT 92, 227–242.
- Ibidhi, R., Frijia, A., Jaouad, M., Salem, H.B., 2018. Typology analysis of sheep production, feeding systems and farmers strategies for livestock watering in Tunisia. Small Ruminant Research 160, 44–53. <https://doi.org/10.1016/j.smallrumres.2018.01.010>.
- International Livestock Research Institute (ILRI), International Union for Conservation of Nature (IUCN), Food and Agriculture Organization (FAO), World Wildlife Fund (WWF), United Nations Environment Program (UNEP) and International Land Coalition (ILC). 2021. Rangelands Atlas. ILRI, Nairobi, Kenya.
- Jaouad, M., Sghaier, M., Bechir, R., Khorchani, T., 2022. Au-delà de la sécurité alimentaire et nutritionnelle: quelle importance de l'élevage pastoral dans les filières d'élevage dans le sud-est tunisien? Revue Européenne du Droit Social 54, 112–120.
- Jayasuriya, S., Mudbhary, P., Broca, S.S., 2012. Food price spikes, increasing volatility and global economic shocks: coping with challenges to food security in Asia. RAP Publications, Food and Agriculture Organization of the United Nations Regional Office for Asia and the Pacific, Bangkok, Thailand.
- Jeder, H., Khalifa, A.B., Sghaier, M., 2013. Impact des changements climatiques sur l'agriculture dans la plaine de Jeffara sud-est tunisien. Journal of Agriculture and Environment for International Development 107, 229–242. <https://doi.org/10.12895/jaeid.20132.164>.
- Jemaa, T., 2016. Stratégies d'adaptation des systèmes d'élevage ovins et modes d'utilisation des parcours en Tunisie Centrale PhD Thesis. Université Montpellier SupAgro, Montpellier, France.
- Jemli, M.H., Boulajfene, H., Azouzi, Z., Ben Salem, W., Khaldi, S., 2018. Projet de développement de l'élevage camelin en Tunisie. Revue Marocaine des Sciences Agronomiques et Vétérinaires 6, 256–329.
- Khosravi Mashizi, A., Heshmati, G.A., Salman Mahini, A.R., Escobedo, F.J., 2019. Exploring management objectives and ecosystem service trade-offs in a semi-arid rangeland basin in southeast Iran. Ecological Indicators 98, 794–803. <https://doi.org/10.1016/j.ecolind.2018.11.065>.
- Lorent, H., Sonnenschein, R., Tsiourlis, G.M., Hostert, P., Lambin, E., 2009. Livestock subsidies and rangeland degradation in central Crete. Ecology and Society 14, 2.
- Maatougui, A.C., 2000. Évolution des systèmes de production au Maroc Occidental Central. Actes de 5ème Séminaire International du Réseau Parcours, 16–18 avril 1998, El Jadida, Morocco, pp. 63–75.
- Mittal, A., 2009. The 2008 food price crisis: rethinking food security policies. G-24 Discussion Paper No. 56. United Nations Publication, New York, NY, USA and Geneva, Switzerland.
- Mohamed, C., Dhoui, A., Ben-Nasr, J., 2021. Economics and Profitability of Goat Breeding in the Maghreb Region. In Goat Science-Environment, Health and Economy (ed. Sándor Kukovics). IntechOpen, London, UK, no. 21.
- Morales, F.D.A.R., Genís, J.M.C., Guerrero, Y.M., 2019. Current status, challenges and the way forward for dairy goat production in Europe. Asian-Australasian Journal of Animal Sciences 32, 1256.
- Moslah, M., Hammadi, M., Khorchani, T., 2004. Productivité de l'élevage camelin dans les parcours du Sud tunisien. Cahiers Options Méditerranéennes 62, 343–347.
- Moslam, M., Megdiche, F., 1989. L'élevage camelin en Tunisie. Séminaire sur la Digestion, la Nutrition et l'Alimentation du Dromadaire, 02/27–01/03/1998, Ouargla, Algérie, pp. 33–36.
- Muñoz-Ulecia, E., Bernués, A., Casasús, I., Olaizola, A.M., Lobón, S., Martín-Collado, D., 2021. Drivers of change in mountain agriculture: A thirty-year analysis of trajectories of evolution of cattle farming systems in the Spanish Pyrenees. Agricultural Systems 186, 102983.
- Nasr, N., 2004. Agriculture et émigration dans les stratégies productives des jbalia du sud-est tunisien. In Environnement et sociétés rurales en mutation. Approches alternatives (ed. Picouet, M., Sghaier, M., Genin, D., Abaab, A., Guillaume, H., Eloumi, M.). IRD éditions, Paris, France, pp. 248–256.
- National Institute of Statistics of Tunisia (INS), 2014. Rapport annuel statistique de la Tunisie de 2018. Retrieved on 8 January 2022 from <http://www.ins.tn/enquetes/recensement-general-de-la-population-et-de-lhabitat-2014>.

- National Institute of Statistics of Tunisia (INS), 2018. Rapport annuel statistique de la Tunisie de 2018. Retrieved on 10 January 2022 from <http://www.ins.tn/publication/tunisie-en-chiffres-2018>.
- Neffati, M., 2020. Capitalisation de l'expérience tunisienne et valorisation des acquis dans le domaine du développement des territoires pastoraux. *Revue des Régions Arides- Numéro Spécial* 47, 2.
- Nefzaoui, A., Ketata, H., El Mourid, M., 2012. Changes in North Africa production systems to meet climate uncertainty and new socio-economic scenarios with a focus on dryland areas. *Options Méditerranéennes: Série A. Séminaires Méditerranéens* 102, 403–421.
- Nefzaoui, A., 2004. Rangeland improvement and management options in arid environment of Central and South Tunisia. In *Réhabilitation des pâturages et des parcours en milieux méditerranéens* (ed. Ferchichi A.). *Cahiers Options Méditerranéennes* 62, 17–26.
- Nefzi, A., 2012. Evaluation économique de l'impact du changement climatique sur l'agriculture: étude théorique et application au cas de la Tunisie PhD Thesis. AgroParisTech, Paris, France.
- Olson, D.M., Dinerstein, E., 2002. The Global 200: Priority ecoregions for global conservation. *Annals of the Missouri Botanical Garden* 89, 199–224.
- Oñatibia, G.R., Aguiar, M.R., Semmartin, M., 2015. Are there any trade-offs between forage provision and the ecosystem service of C and N storage in arid rangelands? *Ecological Engineering* 77, 26–32. <https://doi.org/10.1016/j.ecoeng.2015.01.009>.
- Paniagua, A., 2019. Voices in the new millenium. Shepherds in the Pela Mountains (Southwest of Soria. Central Spain). *Advances in Applied Sociology* 9, 551–568. <https://doi.org/10.4236/aasoci.2019.912040>.
- Powell, J.M., Pearson, R.A., Hiernaux, P.H., 2004. Crop-Livestock Interactions in the West African Drylands. *Agronomy Journal* 96, 469–483. <https://doi.org/10.2134/agronj2004.4690>.
- R Core Team, 2019. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. URL: <https://www.R-project.org/>.
- Ratemo, C.M., Ogendi, G.M., Huang, G., Ondieki, R.N., 2020. Application of traditional ecological knowledge in food and water security in the Semi-Arid Turkana County, Kenya. *Open Journal of Ecology* 10, 321–340.
- REACH and Mercy Corps, 2018. Tunisia, country of emigration and return. Migration dynamics since 2011. REACH, Geneva, Switzerland.
- Richard, J.F., 2006. Le devenir de l'agriculture tunisienne face à la libéralisation des échanges. *Afrique Contemporaine* 219, 29–42.
- Rigolot, C., Martin, G., Dedieu, B., 2019. Renforcer les capacités d'adaptation des systèmes d'élevage de ruminants: Cadres théoriques, leviers d'action et démarche d'accompagnement. *INRAE Productions Animales* 32, 1–12. <https://doi.org/10.20870/productions-animales.2019.32.1.2414>.
- Ryschawy, J., Joannon, A., Choisis, J.P., Gibon, A., Le Gal, P.Y., 2014. Participative assessment of innovative technical scenarios for enhancing sustainability of French mixed crop-livestock farms. *Agricultural Systems* 129, 1–8. <https://doi.org/10.1016/j.agsy.2014.05.004>.
- Selmi, S., Elloumi, M., 2007. Tenure foncière, mode de gestion et stratégies des acteurs. Le cas des parcours du Centre et du Sud tunisien. *Vertigo. La revue électronique en sciences de l'environnement, Hors-série* 4. Retrieved on 21 February 2022 from <http://journals.openedition.org/vertigo/695>.
- Selmi, C., Jaouad, M., Faye, B., Haouat, F., 2018. Typologie des éleveurs camelin au sud-est tunisien en vue de leurs performances économiques. *Colloque International Développement socio-économique et dynamique des sociétés rurales: Pluralité d'acteurs, gestion des ressources et développement territorial*, 3–5 Mai 2016, Zarzis, Tunisie. *Revue des Régions Arides numéro spécial* 44, 209–214.
- South Development Office (ODS), 2018. Retrieved on 10 February 2022 from <http://www.ods.nat.tn/fr/index.php?id=32>.
- Steinfeld, H., Wassenaar, T., Jutzi, S., 2006. Livestock production systems in developing countries: status, drivers, trends. *Revue scientifique et technique* 25, 505–516.
- Trabelsi, M.A., 2016. Analyse de la filière viande rouge en Tunisie: viande cameline. Institut National Agronomique de Tunis, Tunis, Tunisie.
- Valbuena, D., Groot, J.C., Mukalama, J., Gérard, B., Tiftonell, P., 2015. Improving rural livelihoods as a "moving target": trajectories of change in smallholder farming systems of Western Kenya. *Regional Environmental Change* 15, 1395–1407. <https://doi.org/10.1007/s10113-014-0702-0>.
- Vall, E., Marre-Cast, L., Kamgang, H.J., 2017. Chemins d'intensification et durabilité des exploitations de polyculture-élevage en Afrique subsaharienne: contribution de l'association agriculture-élevage. *Cahiers Agricultures* 26, 1–12.
- Vesco, P., Kovacic, M., Mistry, M., Croicu, M., 2021. Climate variability, crop and conflict: Exploring the impacts of spatial concentration in agricultural production. *Journal of Peace Research* 58, 98–113. <https://doi.org/10.1177/0022343320971020>.
- Volpato, G., King, E.G., 2019. From cattle to camels: trajectories of livelihood adaptation and social-ecological resilience in a Kenyan pastoralist community. *Regional Environmental Change* 19, 849–865. <https://doi.org/10.1016/j.livsci.2019.03.017>.