

Applying the ecosystem services framework to pasture-based livestock farming systems in Europe

T. Rodríguez-Ortega¹, E. Oteros-Rozas^{2,3}, R. Ripoll-Bosch^{1,4}, M. Tichit^{5,6}, B. Martín-López³ and A. Bernués^{7†}

¹Centro de Investigación y Tecnología Agroalimentaria de Aragón (CITA). Av. Montañana 930, 50059 Zaragoza, Spain; ²Public Administration and Policy Group, Wageningen University, Hollandseweg 1 6706KN Wageningen, The Netherlands; ³Social-Ecological Systems Laboratory, Department of Ecology, Universidad Autónoma de Madrid, Cantoblanco s/n, 28049 Madrid, Spain; ⁴Animal Production Systems Group, Wageningen University, P.O. Box 338, 6700 AH Wageningen, The Netherlands; ⁵INRA, UMR 1048 SAD APT, F-75231 Paris 05, France; ⁶AgroParisTech, UMR 1048 SAD APT, F-75005 Paris, France; ⁷Department of Animal and Aquacultural Sciences, Norwegian University of Life Sciences (NMBU), 1432 Ås, Norway

(Received 24 September 2013; Accepted 13 February 2014; First published online 12 March 2014)

The concept of 'Ecosystem Services' (ES) focuses on the linkages between ecosystems, including agroecosystems, and human well-being, referring to all the benefits, direct and indirect, that people obtain from ecosystems. In this paper, we review the application of the ES framework to pasture-based livestock farming systems, which allows (1) regulating, supporting and cultural ES to be integrated at the same level with provisioning ES, and (2) the multiple trade-offs and synergies that exist among ES to be considered. Research on livestock farming has focused mostly on provisioning ES (meat, milk and fibre production), despite the fact that provisioning ES strongly depends on regulating and supporting ES for their existence. We first present an inventory of the non-provisioning ES (regulating, supporting and cultural) provided by pasture-based livestock systems in Europe. Next, we review the trade-offs between provisioning and non-provisioning ES at multiple scales and present an overview of the methodologies for assessing biophysical trade-offs. Third, we present non-biophysical (economical and socio-cultural) methodologies and applications for ES valuation. We conclude with some recommendations for policy design.

Keywords: multifunctionality, public goods, biodiversity, valuation, trade-offs

Implications

Pasture-based livestock systems are multifunctional, delivering multiple services to society. Ecosystem Services (ES) are defined as the direct and indirect contributions of ecosystems to human well-being, many of which do not have a market value and are ignored within evaluation frameworks. Designing more informed agro-environmental policy requires us to: explore, identify and evaluate the whole range of ES linked to pasture-based livestock systems; describe the biophysical relationships among different types of agro-pastoral practices; unravel the trade-offs among different types of ES; and value the different types of ES from a three-dimensional (ecological, socio-cultural and economic) perspective.

Introduction

The concept of Ecosystem Services (ES), publicised by the Millennium Ecosystem Assessment (2005), focuses on the linkages between ecosystems, including agroecosystems, and human

well-being. ES are all the contributions, direct and indirect, that people obtain from ecosystems (de Groot *et al.*, 2010). The contributions can be monetary or socio-cultural (i.e. the benefits provided by ecosystems to users' cultural identity, spiritual values or social relationships (Chan *et al.*, 2012a)).

Formally, ES are classified into four groups: *provisioning* ES are material or energy outputs including goods such as food, water, fuel, timber and fibre; *regulating* ES are biophysical processes providing benefits such as climate regulation, flood prevention, waste treatment and water purification; *cultural* ES are recreational, aesthetic and spiritual benefits provided by ecosystems; and *supporting* ES, such as soil formation, photosynthesis or nutrient cycling, are the various processes that are necessary for the production of all the other ES. The *non-provisioning* ES (i.e. the *regulating*, *supporting* and *cultural* ES) mostly constitute public goods; individuals cannot be excluded from their use, and their use by one individual does not reduce their availability to other individuals (Cooper *et al.*, 2009).

Although its connection with the ES framework has been limited, agroecology has several features that are remarkably

† E-mail: alberto.bernues@nmbu.no

similar to those of ES. First, agriculture is considered a multifunctional, social–ecological system, delivering not only marketable goods but also a wide range of public goods. Second, multidimensionality is important in the valuation of agroecosystems and includes biophysical, economic and socio-cultural values. Third, understanding the complexity of agroecosystems requires a trans-disciplinary approach (Dumont *et al.*, 2013). Fourth, synergies between scientific knowledge and traditional knowledge are needed to assess the diverse values of agroecosystems and to achieve social transformation and incidence in policymaking (Oteros-Rozas, 2013).

Although the application of the ES framework to research on farming systems is still in its infancy, it has the potential to integrate the *provisioning* and *non-provisioning* ES at the same level of priority, allowing the multiple trade-offs and synergies that can exist among ES at different scales to be considered. Several high-profile reviews and meta-analyses have called for the development of multiservice approaches (Kareiva *et al.*, 2007); however, the trade-offs and synergies among the different types of ES are still poorly documented (Seppelt *et al.*, 2011; Nieto-Romero *et al.*, 2014). Moreover, most ES valuation studies on agroecosystems have focused on assessing the delivery of one or two ES, ignoring the ecological and social processes underlying the delivery of the complete set of ES (Power, 2010; Nieto-Romero *et al.*, 2014).

Similarly, the consideration of the full range of ES in pasture-based livestock farming systems (PLFS) is also new, despite the fact that PLFS, more than other agroecosystems, strongly depend on and influence the *regulating* and *supporting* ES (Zhang *et al.*, 2007). PLFS have the potential to ensure multiple ES (e.g. hedgerows reduce erosion, grasslands filter runoff, natural predators control pests and wild bees provide pollination), but there are trade-offs. Managing the trade-offs and synergies among ES at multiple scales is essential for reinforcing the contribution of PLFS to landscape multifunctionality and human well-being. Supporting methodologies that help stakeholders and policymakers to better understand the trade-offs and synergies among ES are needed to help design alternatives and explore scenarios for the future.

The ES approach is stimulating debate about the need to introduce deep policy changes (European Commission, 2011b; Bateman *et al.*, 2013) that integrate agricultural policies with policies in other sectors, such as biodiversity (European Commission, 2011a). It also shifts the emphasis towards the supply of public, non-market goods, and thus opens up opportunities for Payments for Ecosystem Services, or 'green payments'.

The objective of our review was to explore the application of the ES framework to European PLFS. We first established a biophysical inventory of the *non-provisioning* ES provided by PLFS. Second, we focused on the trade-offs between *provisioning* and *non-provisioning* ES at the field, farm, and landscape levels, and the methodologies used for assessing them in biophysical terms. Third, we reviewed the non-biophysical

methodologies for ES valuation and their application to PLFS. Finally, we proposed some recommendations for policy design.

An inventory of the ES provided by PLFS in Europe

The consideration of the *regulating*, *supporting* and *cultural* ES delivered by agroecosystems is still relatively new (Swinton *et al.*, 2007). The biophysical relationships between agroecosystems and the *non-provisioning* ES are neither readily apparent nor easy to measure; however, they provide the knowledge base for their valuation. We reviewed the scientific literature acknowledging those relationships within the context of PLFS. The ES were classified following the definitions given in the Millennium Ecosystem Assessment (2005) and the classification proposed by The Economics of Ecosystems and Biodiversity (Kumar, 2010). Biodiversity is considered a key element supporting the delivery of other ES in both the classification systems and was therefore included in our search.

We conducted a literature search on 11 December 2013 and included all peer-reviewed publications in the SCOPUS database. Our preliminary search included a range of terms related to PLFS and the diverse *non-provisioning* ES (including biodiversity) and resulted in more than 13 500 references. To circumscribe the search within the ES framework, we restricted the query by explicitly including the term 'ecosystem service' and equivalent terms (i.e. 'public good', 'externality', 'environmental service' and 'multifunctionality'). Detailed information regarding the terms included in the query is available in the Supplementary material S1. The final search result included 563 articles and showed a rapid increase in the number of publications (from an average of 10.5 per year in the period 1995 to 2005 up to 228 in 2013) following the appearance of the Millennium Ecosystem Assessment (2005) and the formal establishment of the ES concept (Figure 1). The rates of publication on *supporting* and *cultural* ES increased similarly, but to a lesser degree than the rate of publication on *regulating* ES.

The coverage of the different *non-provisioning* ES in the literature was irregular: 'gene-pool protection' (which includes biodiversity) accounted for the largest share of the publications (30.5%), followed by 'aesthetic value' of landscape (27.3%) and 'climate regulation' (12%); the prevention or moderation of natural hazards (such as 'forest fires' (0.3%)), 'air purification' (0.3%), 'cognitive development' (0.2%) and 'spiritual experience' (0.1%) were studied little in relation to PLFS (Figure 2).

The search results were checked against the following criteria: the PLFS was located in Europe; the study included biophysical quantification; the paper described original research; the experimental design and analysis were sound; and the results showed relations between land management or agricultural practices and ES. Thirty publications met all of the criteria for inclusion in our subsequent analysis (Supplementary material S2). Several reasons led to the dismissal of most publications: studies were conducted in crop/arable land or forestry areas lacking explicit links to

pasture-based systems; relationships between livestock and ES were mentioned but not explicitly assessed; the ES were not assessed in biophysical terms (as in reviews, theoretical reflections or frameworks for analysis) or followed non-biophysical methodologies (e.g. economic valuation or policy analysis).

We classified the 30 references meeting our criteria according to the type of ES and the indicators used in the assessment, the geographic scale of the study (patch/field, farm, or region/landscape) and the factor under analysis (e.g. land-use change, management regime or agricultural practice; Table 1). The quantitative methodologies and the main evidences found are listed in Supplementary Table S1.

Most of the studies meeting our criteria focused on biodiversity and landscape. Despite the different approaches and methodologies, there was general agreement that large shifts in PLFS management tend to impair biodiversity (Chamberlain *et al.*, 2000; Stoate *et al.*, 2001). Biodiversity was negatively affected by landscape homogenisation, either because of agricultural intensification or abandonment (Plieninger *et al.*, 2006), and positively affected by the use of low-input, heterogeneous and restored PLFS (Weigelt *et al.*,




Figure 1 The number of publications per year and the type of ecosystem services covered (see the Supplementary material S1 for query specifications). Biodiversity is included in the supporting ecosystem services.

2009; Albrecht *et al.*, 2010; Varah *et al.*, 2013). Grasslands in Europe, as man-made habitats, need to be managed for higher structural heterogeneity to maintain high species diversity (Diacon-Bolli *et al.*, 2012).

Landscape was analysed from diverse perspectives: vegetation dynamics, landscape diversity or aesthetic quality. In terms of vegetation change, we found reports of a general process of abandonment in Europe (e.g. MacDonald *et al.*, 2000), especially in less favoured areas (mountainous, less productive or marginal areas) where PLFS have declined (Bernués *et al.*, 2005). This abandonment has led to a general trend of afforestation and encroachment, ending in landscape closure or homogenisation (Tasser *et al.*, 2007; Riedel *et al.*, 2013), reducing the mosaic value (Brady *et al.*, 2012). Moreover, afforestation impacts community composition (Alrababah *et al.*, 2007; Buscardo *et al.*, 2008); rare or specialist species are often replaced by habitat generalists or species associated with forested habitats (Oxbrough *et al.*, 2006).

Although aesthetic preferences are highly subjective and incorporate social constructs, some predictive variables were proxies for attractiveness to society, such as the presence of water, the number of different land types and the heterogeneity of the landscape (Dramstad *et al.*, 2006). In this sense, Ford *et al.* (2012) concluded that environmental appreciation would be greater in grazed areas, because those areas sustain significantly greater plant species richness, particularly among forbs flowers. Garcia-Llorente *et al.* (2012) found that multifunctional grasslands delivered not only greater aesthetic value but also a more diverse flow of ES.

Carbon sequestration was usually analysed at the field/patch scale, but direct relationships between it and PLFS were difficult to establish. Carbon stocks were greater under pastures than under cropland (Lorencová *et al.*, 2013), and




Figure 2 The number of publications per ecosystem service (see the Supplementary material S1 for query specifications). Biodiversity is included in the gene pool protection service.

Note: publications that covered more than one ecosystem service are counted more than once.

Table 1 The number of times the influences on ecosystem services of diverse land management and agricultural practices were measured at different spatial scales (n = 30)

	Land-use changes (region/landscape)				Farming system (farm)			Agricultural practices (field/patch)					Total
	Intensification	Abandonment/ encroachment	Land-use types	Other	Intensification	Land-use types	Conventional v. organic	Intensity of management	Abandonment of practice	Land-use type	Stocking rate	Other	
Regulating													
Carbon sequestration			1							1	2	3 ^{c,d,f}	7
Pollination	1					1					1		3
Soil erosion			1										1
Flood control											1		1
Supporting													
Biodiversity		1	1	2 ^a	3	2	3	1		2	4	2 ^{f,g}	19
Nutrient cycling									2		1	1 ^e	2
Cultural													
Landscape	1	3		2 ^{a,b}							1		6
Traditional ecological knowledge	1	1											1
Total	2	3	2	3	3	2	3	1	2	3	6	5	

^aCAP measures, abandonment and intensification.^bStocking rates.^cFertiliser application and plant seeding.^dNumber of species/varieties cultivated.^eManure application, mowing/unmowing.^fNutrient addition (various combinations of mineral nutrients (N, P, K, Mg), lime addition and herbivore exclusion (insect, rabbit)).^gPrescribed burning.

grazing did not compromise the carbon-storage potential (Medina-Roldán *et al.*, 2012). Carbon storage was strongly modulated, however, by location, soil type (Marriott *et al.*, 2010), stocking rate (Martinsen *et al.*, 2011) and grassland management (Soussana *et al.*, 2004).

Many ES did not appear in the studies included in our final search results, probably because the environmental effects (e.g. soil fertility, soil erosion and waste treatment) of PLFS have been studied without any explicit reference to ES. Other ES were important only in particular regions; wild forest fires, for example, have enormous environmental, social and political relevance in Euro-Mediterranean regions but less in northern areas. Grazing by domestic livestock reduced the wildfire risk by limiting the shrub and herbage biomass and maintaining landscape heterogeneity (Ruiz-Mirazo and Robles, 2012); however, the stocking rates and grazing regimes in many cases were not enough to avoid vegetation dynamics towards encroachment (Riedel *et al.*, 2013).

The ES concept is highly complex, because it connects ecological and social systems and recognises particular ES that are strongly connected to biodiversity (i.e. *supporting* and *regulating* ES) or to humans (i.e. *cultural* ES). Despite a diverse spectrum of disciplines is required to study this complexity (Martín-López *et al.*, 2014), we have found that the scope of the biophysical studies differed widely and lacked standardised methodologies. Furthermore, the spatial and temporal scales were often not identified.

The agricultural practices and management regimes under consideration were also very diverse. On the one hand, the studies involved or compared different land-cover types, such as grasslands, woodlands, crops, or fallow or abandoned lands (e.g. Dallimer *et al.*, 2010; McMahon *et al.*, 2010). On the other hand, the livestock management practices were widely heterogeneous, involving the maintenance, or the implementation, of one or several changes in existing practices, such as mowing, grazing and fertilising. For example, some studies compared different grazing intensities (stocking rates) or season lengths (e.g. Batáry *et al.*, 2010; Martinsen *et al.*, 2011), or tested differences between grazed and non-grazed areas (e.g. Medina-Roldán *et al.*, 2012, Riedel *et al.*, 2013), combinations of grazed, mowed and abandoned areas (e.g. Franzén and Nilsson, 2008; Hoiss *et al.*, 2013); or combinations of grazing intensities and fertilisation regimes (e.g. Marriott *et al.*, 2010; Michaud *et al.*, 2012). All of the management regimes had broad effects on ES delivery because of the complex interrelationships among land uses, biodiversity and ES. Hence, establishing comparisons between studies was not straightforward.

Trade-offs among ES at the field, farm and landscape levels

The simultaneous production and the complex, dynamic interrelatedness of multiple ES is often overlooked (Bennett *et al.*, 2009). An overly narrow focus on maximising a limited set of ES could lead to unexpected trade-offs or to undesirable and sudden declines in other ES (Bennett *et al.*, 2009).

Trade-offs among ES occur when the delivery of one service is reduced as a consequence of the increased use of another service. In contrast, synergies among ES arise when multiple ES are enhanced simultaneously (Raudsepp-Hearne *et al.*, 2010). The limited knowledge of the biophysical relationships among ES makes it difficult not only to predict trade-offs and synergies but also to understand the mechanisms that cause them, and hence how to minimise or enhance them. The simultaneous management of multiple ES is important; however, it is also extremely challenging. A better understanding of the processes by which agricultural practices and management regimes influence trade-offs and synergies among ES would allow the outputs of a range of ES to be envisioned (Power, 2010).

Trade-offs across multiple scales

A few recent studies have explored trade-offs within PLFS at multiple spatial scales (e.g. Tichit *et al.*, 2007; Sabatier *et al.*, 2012; Sabatier *et al.*, 2013). These studies used a process-based modelling approach that provides a framework for assessing trade-offs on multiple scales (field, farm and landscape) in grassland-dominated landscapes, where livestock grazing may conflict with bird conservation. The approach integrates the two ecological processes, nest trampling and chick survival, through which the direct and indirect effects of grazing and mowing influenced the life cycles, and thus the population dynamics, of birds. Nest trampling by livestock has a direct, negative effect on bird fecundity. Grass height can have a positive, indirect effect on chick survival. The modelling approach was used to find out how grasslands should be managed to reconcile fodder production, a *provisioning* ES, and bird conservation, a *cultural* ES.

At the field level, Sabatier *et al.* (2010) showed that increased grazing intensity did not have the same impact on all bird species: some species were more sensitive to nest trampling than others. It was necessary to fine-tune the grazing intensity over time to ensure the conservation of several bird species. The tuning involved temporal shifts in grazing sequences to minimise nest trampling and create optimal grass heights for bird survival (Tichit *et al.*, 2005a and 2005b; Durant *et al.*, 2008). The temporal shifts were action levers in the trade-off between the ES, because they improved bird conservation without causing major loss of fodder production.

At the farm level, different land uses, such as mowing and grazing, offered contrasting habitats for birds and contrasting feeding resources for livestock (Martin *et al.*, 2009). The interactions between the ES determined both the herbage production and the health of the bird populations (Sabatier *et al.*, 2010; Tichit *et al.*, 2011). Mowing and intensive grazing increased herbage production but impaired bird fecundity and chick survival. The proportions of grazed/mown fields in the farm area thus formed the key action lever for modulating the supplies of both ES. The study stressed, however, that it was less costly to implement the action lever on extensive (<1.4 livestock units/ha) farms than on intensive farms (>1.4 livestock units/ha) (Tichit *et al.*, 2011).

Similar results have been reported in mountain PLFS, where the level of farm intensification was identified as a key variable modulating the trade-off between cattle production and grassland floristic composition (Jouven and Baumont, 2008).

At the landscape scale, land use was an important factor influencing the provision of, and the relationships among, multiple ES (Foley *et al.*, 2005). It is important to determine whether land-use intensity and allocation can be action levers to move or modify the shape of the trade-off frontier. Recent studies highlighted landscape heterogeneity, the spatial arrangement of the different land uses, as a factor promoting the diversity of available habitats and thus allowing biodiversity to increase (Haslem and Bennett, 2008; Groot *et al.*, 2010). Sabatier *et al.* (2013) went a step further by demonstrating that heterogeneous land use in grassland was an efficient lever to move the trade-off frontier. The simulation of a large number of landscapes revealed that at a given level of *provisioning* service (herbage production), increasing the landscape heterogeneity could improve *cultural services* (bird populations) by changing the spatial arrangement of mowed and grazed areas. The benefits of heterogeneous land use emerged from a set of interacting suboptimal habitats, where each type of land use provided some of the resources needed, and species mobility among land-use types enabled the populations to obtain all the needed resources. Those results were consistent with the results of other modelling studies conducted at the landscape scale (Polasky *et al.*, 2005; Groot *et al.*, 2007), illustrating how strategic land-use placement can improve trade-offs.

Methods for exploring trade-offs

There are several approaches to analysing trade-offs among ES (Groot *et al.*, 2009). Here, we present a short overview of three types of modelling approaches: pareto-based optimisation, co-viability analysis (CVA) and companion modelling. The approaches offer different options to assess multiple ES, quantify the relationships among ES and explore the range of potential solutions.

The common point is to analyse the set of ES as a problem of multi-criteria decision making, where several antagonist criteria are optimised simultaneously by modifying a common driver (e.g. the land-use intensity and its spatial allocation) of several ES.

Pareto-based multi-criteria optimisation offers a static framework to solve spatially explicit problems and find the optimal spatial allocation of land-use intensity. Optimal allocations are those providing the best reconciliation among the different ES. A Pareto frontier graphically represents a set of non-dominated solutions such that a given service could not be improved without deteriorating another service and vice versa (Groot *et al.*, 2012).

CVA provides a quantitative tool for dynamically exploring a solution space defined by the supply of several ES. With CVA, a desired future and road to it can be defined by a set of restrictions representing the limits within which the supply of each ES should be maintained (Tichit *et al.*, 2007, Sabatier *et al.*, 2010 and 2012). Different mathematical tools are used

to compute the set of viable decisions leading to the desired future. This approach is particularly useful for involving stakeholders in a negotiation-based planning process, because restrictions on ES can be set at different levels, depending on the knowledge and priorities of the stakeholders. Furthermore, CVA offers an integrated criterion for achieving multiple goals in a short- to long-term perspective. CVA is a powerful tool for examining interactions across temporal scales and the ways in which different objectives may conflict in the long term as a consequence of short-term decisions.

The boundaries of what is possible in terms of ES delivery should not be limited to mathematical exploration, mainly because one of the underlying aspects of trade-offs is that different stakeholders pursue different, sometimes antagonist, goals on a given landscape. As a consequence, they need to develop a common view on problem and collectively design solutions. Companion modelling is a participatory methodology that involves stakeholders in the different steps of exploring and designing solutions (Souchère *et al.*, 2010). It provides stakeholders with elements for reflection, helping to reinforce planning for the future and understanding of complex multi-scale problems. They act as tools for decision support, dialogue and communication among a variety of stakeholders pursuing multiple objectives. Such tools can account for scale mismatch (Cumming *et al.*, 2006) between ecological processes and human processes, when, for instance, gains emerge on one scale and costs are supported on another. Participatory tools can incorporate scenarios and public policy instruments that are unaccounted for in most studies (e.g. Seppelt *et al.* (2011) found that only 29% of studies accounted for such instruments). We have stressed that it is necessary to account for both the technical and the social and economic dimensions of ES-management practices. Companion modelling offers the advantage of integrated assessment tools that include descriptions of biophysical processes, the influences of individual and collective management decisions on those processes, the perceptions of stakeholders regarding the environment, and the social and economic consequences of management. Thus, it has the potential to foster knowledge and social representation sharing of ES. Through scenario exploration, it can also simulate the impact of management changes on ES delivery. As tools conceived for helping stakeholders to organise themselves and consult each other, they can facilitate the design of new spatial layouts for livestock farming systems that provide diversified mosaics of ES in agroecosystems.

Uncovering the socio-cultural and economic values of ES

Uncovering ES values for human well-being requires diverse tools that embrace the multidimensional (i.e. biophysical, socio-cultural and economic) nature of the value of ES (Martín-López *et al.*, 2014). The plurality of values makes ES intrinsically incommensurable and therefore impossible to reduce to a single unique measure (Gómez-Baggethun and

de Groot, 2008). This is one reason, among others, why different types of ES-valuation methods have been developed.

Socio-cultural valuation of ES

Socio-cultural assessments focus on the preferences, needs, values, norms and behaviours of individuals, institutions and organisations towards ES (Cowling *et al.*, 2008). These valuation approaches are particularly appealing because of their suitability for uncovering the motivations for conserving ES (Chan *et al.*, 2012b). Socio-cultural valuation methods have been praised for their suitability and sensitivity in assessing PLFS (Oteros-Rozas *et al.*, 2013a), because PLFS have been shaped by long-term human activities and therefore have particular cultural values (Martín-López *et al.*, 2012).

The main tools or methods for socio-cultural valuation are consultative methods (structured processes of inquiry into people's perceptions and preferences) and deliberative and participatory methods (group-based activities to elucidate people's relationships with ecosystems, identify conflicts between the beneficiaries of ES, and identify trade-offs between different management strategies, land uses or possible future scenarios) (de Groot *et al.*, 2010; Christie *et al.*, 2012).

Consultative methods include individual questionnaires and in-depth interviews; both tools allow for qualitative and quantitative data analyses, but the questionnaires tend to focus on gathering quantitative data (Struhsaker *et al.*, 2005), whereas the in-depth interviews are more suitable for collecting qualitative data. Deliberative and participatory approaches include focus groups, Delphi surveys, participatory rural appraisal and participatory scenario planning (for a review see Christie *et al.*, 2012). These methods intend to elucidate information about people's relationships with livestock systems to reach consensus or to unravel disagreements about relationships, identify conflicts among the beneficiaries of ES, and identify trade-offs between different management strategies, land uses or possible future scenarios (e.g. Bernués *et al.*, 2013). Interest in participatory mapping of ES through social elicitation has been growing recently (e.g. Palomo *et al.*, 2013), particularly for the spatially explicit quantification of *cultural* ES (Plieninger *et al.*, 2013).

Despite the recognised multifunctionality of agroecosystems, and particularly that of PLFS, there are few examples of socio-cultural valuation of ES related to livestock farming systems. Davies and Hatfield (2007) reviewed the direct and indirect value of ES provided by pastoralism in eight regional studies and highlighted the gaps in knowledge and policy options to support rangeland economies. Oteros-Rozas *et al.* (2012 and 2013b) identified the ES related to transhumance in Spain by carrying out socio-cultural assessments via interviews, cognitive and visual ES-perception surveys, focus groups and participatory scenario planning. Among the 34 ES they identified, the most important for social well-being were fire prevention, air purification and livestock production. They also assessed the trends and factors affecting ES flows and the link between ES and the practice of transhumance on

foot and found that the delivery of certain *regulating* ES (tree regeneration, seed dispersal, and the maintenance of soil fertility and connectivity) and *cultural* ES (local ecological knowledge, cultural exchange and cultural identity) were closely related to transhumance. The results suggested a particularly close link between fire prevention and the maintenance of transhumance in the study area. Lamarque *et al.* (2011) found that different stakeholders appreciated ES differently in three mountainous regions in Europe primarily used for livestock farming. They identified, however, a common set of ES that all stakeholders considered important, including soil erosion, water quantity and quality, forage quality, conservation of plant diversity, and aesthetics and recreation. Pereira *et al.* (2005) used a range of tools including participatory rural appraisal and other field methods to socio-culturally assess the ES provided by agro-pastoral ecosystems in the rural community of Sistelo in northern Portugal. They identified an emotional attachment to livestock and pastoral practices, which, together with a relationship between cattle and the maintenance of pasturelands, hindered encroachment caused by natural succession, thus preventing wildfires. Heikkinen *et al.* (2012) elaborated on the relationship between pastoralism and ES from the user and producer perspectives through an exercise in scenario analysis. They modelled biodiversity conservation and ecotourism as ES produced by pastoral communities under scenarios representing novel solutions to conservation–pastoralism dilemmas. Bernués *et al.* (2013) used focus groups to quantify the importance that farmers and non-farmers attached to the ES delivered by mountain agriculture. They found that aesthetics (landscape/vegetation), gene pool protection (biodiversity maintenance) and natural hazard prevention (forest fires) were, together with opportunities for recreation and culture, were the most important ES delivered by mountain livestock systems. Several of the previously mentioned studies indicated differences between the perceptions of farmers and non-farmers: farmers gave more importance to *regulating* and *provisioning* ES, mainly those related to their own farming activity or local circumstances; whereas non-farmers gave more importance to *cultural* ES, generally showing more global concerns.

Economic valuation of ES

The economic valuation of ES can provide useful information about the monetary gains and losses caused by different land-use management options; thus, it can be a useful tool to quantify the ES trade-offs among different management options (Hicks *et al.*, 2009; Martín-López *et al.*, 2011). PLFS have great economic importance, namely, their total economic value (TEV), despite the fact that conventional markets do not recognise most ES. Components of the TEV are usually represented by a value taxonomy, which distinguishes between *use* and *non-use* values (Table 2).

On the one hand, *use* values comprise *direct use*, *indirect use* and *option* values. *Direct use* values derive from the conscious use and enjoyment of ES. They may be *extractive*, such as food, or *non-extractive*, such as recreation, nature

Table 2 Relationships between the components of total economic value (TEV) and the ecosystem services delivered by pasture-based farming systems

	Component of TEV	Ecosystem services category	Particular ecosystem services	Proper economic valuation technique
Less difficult to estimate	Use values			
	Direct-use value			
	Extractive	Provisioning	Food, timber, fibre, hunting	Direct market analysis and production function analysis
	Non-extractive	Cultural	Recreation and tourism Aesthetic enjoyment	Travel-cost method Hedonic pricing Contingent valuation Choice experiment
	Indirect-use value	Regulating	All	Avoided or replacement costs Contingent valuation Choice experiment
	Option value	All	All	Contingent valuation Choice experiment
More difficult to estimate	Non-use values			
	Bequest value	All	All	Contingent valuation Choice experiment
	Existence value	Cultural	Satisfaction for conserving agrobiodiversity	Contingent valuation Choice experiment

The most suitable economic valuation techniques for each of the TEV components and ecosystem services are shown.

Table 3 Characterisation of the main economic valuation approaches used for placing monetary value on ecosystem services

Economic valuation approach	Characteristics	Specific techniques	Application to pasture-based livestock farming systems
Market-based	Based upon current markets	Direct market analysis Production function analysis Replacement or avoided costs	Scoones (1992), Oteros-Rozas <i>et al.</i> (2012)
Revealed-preference	Infers values from human behavioural changes in real markets that are related in some way to the ecosystem service	Travel cost Hedonic pricing	Shonkwiler and Englin (2005) Ready and Abdalla (2005), Ma and Swinton (2011)
Stated-preference	Estimates economic values through hypothetical markets	Contingent valuation Choice modelling	Scarpa <i>et al.</i> (2003), Zander and Drucker (2008), Ruto <i>et al.</i> (2008) Zander <i>et al.</i> (2013)
Benefit transfer	Infers the value of ecosystem services based on previous estimations at another study site	Unit value Benefit function Meta-analysis function transfer	Not explicitly applied to livestock farming system

tourism and aesthetics enjoyment. Hence, *extractive direct use* values are strongly related to *provisioning* ES, and *non-extractive direct use* values are related to *cultural* ES. *Indirect use* values are associated with the *regulating* ES delivered by livestock farming systems and do not entail conscious enjoyment or use. Finally, *option* values are related to future direct and indirect uses by humans. *Non-use* values are those arising from people’s feelings towards the existence of biodiversity and the ES that biodiversity provides. This type of value includes the satisfaction of knowing that ES will be available to future generations (*bequest* value) and the satisfaction of knowing that species, farmlands or ES continue to exist (*existence* value). Because *non-use* values are strongly based on moral and ethical issues concerning

future human generations and biodiversity, they are extremely difficult to estimate with any precision.

The broad spectrum of economic valuation methods that exist to cope with the heterogeneous values derived from ES can be classified into four basic approaches: market, revealed-preference, stated-preference and benefit transfer (Table 3). These approaches (except for benefit transfer) estimate the monetary value of ES on the basis of stakeholders’ preferences, expressed either in real markets (market and revealed-preference methods) or in hypothetical markets (stated-preference methods).

The broad landscape of economic techniques allows us to estimate the monetary value of the ES provided by PLFS (Table 3). Direct market analysis estimates the economic

value of the *provisioning* ES, because most *provisioning* ES (e.g. meat, milk and fibre) have real market prices. The monetary values of ES that do not have markets can be estimated in an indirect way by analysing related markets using (1) the costs avoided because the maintenance of certain *regulating* ES, such as soil fertility (e.g. the manure produced by sheep allowed farmers to avoid the costs of fertilisation (Oteros-Rozas *et al.*, 2012)); (2) the travel-cost method for assessing the *cultural* ES of recreational activities and nature tourism (e.g. Shonkwiler and Englin, 2005; Pouta and Ovaskainen, 2006); and (3) the hedonic pricing for *cultural* ES, such as the recreational or aesthetic values derived from farmlands (e.g. Ma and Swinton, 2011). In addition, the stated-preference methods provide an alternative, comparative way to estimate the monetary value of most ES. On the one hand, such methods are able to estimate the economic value associated with the most intangible ES, such as the value of the existence of biodiversity (Venkatachalam, 2004). On the other hand, the economic value of different ES can be estimated in the same exercise, because stated-preference methods are performed on the basis of hypothetical markets created through questionnaires in which people state their willingness to pay for preserving ES. For instance, the choice-experiment technique allows us to jointly estimate the economic values of the different ES provided by PLFS (see Table 2). Recent studies have focused on estimating the economic losses associated with the erosion of agro-biodiversity, in terms of diminishing local breeds, through the consideration of different ES, different components of the TEV, and both private and public goods. Martin-Collado (2013) and Zander *et al.* (2013) analysed the social importance of one *provisioning* ES (food quality), one *supporting* ES (gene-pool maintenance) and three *cultural* ES (landscape aesthetics, cultural identity and the existence of the local breeds) attached to three local breeds in Spain and Italy. Their results showed that society valued the non-market ES (public goods) highly, as more than 75% of the estimated economic value arose from the *cultural* ES and the maintenance of the gene pool. Similarly, Kassie *et al.* (2009) found that among the attributes of indigenous cows in central Ethiopia, those indirectly related to markets (i.e. fertility or disease resistance) were as important as those directly related to markets (i.e. milk provisioning). Those results suggest that society highly values the public goods derived from local breeds, suggesting that the associated ES should not be hidden in the decision-making processes of agro-biodiversity management.

Implications for policy design

Some considerations for agro-environmental policy design in Europe can be underlined on the basis of this review.

Incorporate non-provisioning ES into decision making

Agricultural policy design for PLFS should not focus only on *provisioning* ES, because this can result in decisions that reduce the TEV of the system (Bateman *et al.*, 2013).

Because many ES (e.g. spiritual values, cognitive development, and certain *regulating* and *supporting* services) cannot be readily translated into monetary values, or because doing so can be undesirable (Gómez-Baggethun and Ruiz-Pérez, 2011), the socio-cultural values of ES need to be considered across agricultural, environmental and rural development policies.

A few ES should be chosen based on existing scientific evidence and prioritised across different agroecosystems and regions. However, the socio-cultural, economic and biophysical contexts across different sites in Europe strongly influence the valuation of ES (e.g. the prevention of forest fires is key in Mediterranean countries but not in northern Alpine areas), and therefore comparing different case studies and scaling up results can be difficult (Seppelt *et al.*, 2011). Similarly, a small number of agricultural practices and land-management regimes, those with the greatest potential to enhance the prioritised ES, should be targeted by agro-environmental policies.

Consider the trade-offs and synergies

There are trade-offs within PLFS between *provisioning* and *non-provisioning* ES, and between *non-provisioning* services and disservices (negative environmental impacts), but there are also synergies (Bernués *et al.*, 2011). Therefore, European policies should promote farming practices that constitute action levers to maintain a diverse spectrum of ES that benefit different stakeholders.

Individualise compensation schemes and select relevant indicators for monitoring

For agro-environmental measures to constitute effective payments for ES, schemes need to be regionalised and, if possible, individualised by farmers or farmer groups. In addition, because targeted schemes bear closely on local and small farmers' interests, payments for ES should be promoted as rewards rather than subsidies (Sabatier *et al.*, 2012). To do so, we must establish objective, easy-to-understand, measurable and responsive indicators to monitor the effects of compensation schemes on ES delivery and the well-being of farmers.

Finally, in terms of European research policy, *dynamic and multi-scale approaches* for assessing ES are greatly needed because of the mismatches between the temporal and spatial scales of various ES and agricultural practices (e.g. the recurrent labour costs of grazing are short term and occur at the farm level, whereas the benefits of forest-fire prevention are long term and reach wider territories and more recipients). While difficult to perform, upscaling exercises are necessary to contribute to agro-environmental decision making at European scale.

The ES framework integrates the capacity of agroecosystems to supply diverse ES and the perspectives and interests of stakeholders regarding the use of ES. *Inter-disciplinary frameworks* involving natural and social scientists are therefore extremely important. Because the ES concept bears on agroecosystem management and policy, they should be opened not only to scientists and decision makers but also to other stakeholders.

Acknowledgements

Projects: Marie Curie IEF Contract no. 299794 and INIA RTA2011-00133-C02-02. The first author has a pre-doctoral fellowship from INIA.

Supplementary material

To view supplementary material for this article, please visit <http://dx.doi.org/10.1017/S1751731114000421>.

References

- Albrecht M, Schmid B, Obrist MK, Schüpbach B, Kleijn D and Duelli P 2010. Effects of ecological compensation meadows on arthropod diversity in adjacent intensively managed grassland. *Biological Conservation* 143, 642–649.
- Alrababah MA, Alhamad MA, Suwaileh A and Al-Gharaibeh M 2007. Biodiversity of semi-arid Mediterranean grasslands: impact of grazing and afforestation. *Applied Vegetation Science* 10, 257–264.
- Batáry P, Báldi A, Sárospataki M, Kohler F, Verhulst J, Knop E, Herzog F and Kleijn D 2010. Effect of conservation management on bees and insect-pollinated grassland plant communities in three European countries. *Agriculture, Ecosystems and Environment* 136, 35–39.
- Bateman JJ, Harwood AR, Mace GM, Watson RT, Abson DJ, Andrews B, Binner A, Crowe A, Day BH, Dugdale S, Fezzi C, Foden J, Hadley D, Haines-Young R, Hulme M, Kontoleon A, Lovett AA, Munday P, Pascual U, Paterson J, Perino G, Sen A, Siriwardena G, van Soest D and Termansen M 2013. Bringing ecosystem services into economic decision-making: land use in the United Kingdom. *Science* 341, 45–50.
- Bennett EM, Peterson GD and Gordon LJ 2009. Understanding relationships among multiple ecosystem services. *Ecology Letters* 12, 1394–1404.
- Bernués A, Rodríguez-Ortega T, Ripoll-Bosch R and Casasús I 2013. A qualitative research on Spanish farmers and citizens perceptions of ecosystem services provided by mountain livestock farming. In 17th Meeting of the FAO-CIHEAM Mountain Pastures Network, Pastoralism and ecosystem conservation, 5 to 7 June 2013, Trivero, Italy, p. 4.
- Bernués A, Ruiz R, Olaizola A, Villalba D and Casasús I 2011. Sustainability of pasture-based livestock farming systems in the European Mediterranean context: synergies and trade-offs. *Livestock Science* 139, 44–57.
- Bernués A, Riedel JL, Asensio MA, Blanco M, Sanz A, Revilla R and Casasús I 2005. An integrated approach to studying the role of grazing livestock systems in the conservation of rangelands in a protected natural park (Sierra de Guara, Spain). *Livestock Production Science* 96, 75–85.
- Brady M, Sahrbacher C, Kellermann K and Happe K 2012. An agent-based approach to modeling impacts of agricultural policy on land use, biodiversity and ecosystem services. *Landscape Ecology* 27, 1363–1381.
- Buscardo E, Smith GF, Kelly DL, Freitas H, Iremonger S, Mitchell FJG, O'Donoghue S and McKee AM 2008. The early effects of afforestation on biodiversity of grasslands in Ireland. *Biodiversity and Conservation* 17, 1057–1072.
- Chamberlain DE, Fuller RJ, Bunce RG, Duckworth JC and Shrubbs M 2000. Changes in the abundance of farmland birds in relation to the timing of agricultural intensification in England and Wales. *Journal of Applied Ecology* 37, 771–788.
- Chan KMA, Satterfield T and Goldstein J 2012a. Rethinking ecosystem services to better address and navigate cultural values. *Ecological Economics* 74, 8–18.
- Chan KMA, Guerry AD, Balvanera P, Klain S, Satterfield T, Basurto X, Bostrom A, Chuenpagdee R, Gould R, Halpern BS, Hannahs N, Levine J, Norton B, Ruckelshaus M, Russell R, Tam J and Woodside U 2012b. Where are cultural and social in ecosystem services? A framework for constructive engagement. *BioScience* 62, 744–756.
- Christie M, Fazey I, Cooper R, Hyde T and Kenter JO 2012. An evaluation of monetary and non-monetary techniques for assessing the importance of biodiversity and ecosystem services to people in countries with developing economies. *Ecological Economics* 83, 67–78.
- Cooper T, Hart K and Baldock D 2009. Provision of Public Goods through Agriculture in the European Union (Report prepared for DG Agriculture and Rural Development, Contract No 30-CE-0233091/00-28). Institute for European Environmental Policy, London, United Kingdom.
- Cowling RM, Ego B, Knight AT, O'Farrell PJ, Reyers B, Rouget M, Roux DJ, Welz A and Wilhelm-Rechman A 2008. An operational model for mainstreaming ecosystem services for implementation. *Proceedings of the National Academy of Sciences of the United States of America* 105, 9483–9488.
- Cumming GS, Cumming DHM and Redman CL 2006. Scale mismatches in social-ecological systems: causes, consequences, and solutions. *Ecology and Society* 11, 4.
- Dallimer M, Marini L, Skinner AMJ, Hanley N, Armsworth PR and Gaston KJ 2010. Agricultural land-use in the surrounding landscape affects moorland bird diversity. *Agriculture, Ecosystems and Environment* 139, 578–583.
- Davies J and Hatfield R 2007. The economics of mobile pastoralism: a global summary. *Nomadic Peoples* 11, 91–116.
- de Groot RS, Alkemade R, Braat L, Hein L and Willemsen L 2010. Challenges in integrating the concept of ecosystem services and values in landscape planning, management and decision making. *Ecological Complexity* 7, 260–272.
- Diacon-Bolli J, Dalang T, Holderegger R and Bürgi M 2012. Heterogeneity fosters biodiversity: linking history and ecology of dry calcareous grasslands. *Basic and Applied Ecology* 13, 641–653.
- Dramstad WE, Tveit MS, Fjellstad WJ and Fry GLA 2006. Relationships between visual landscape preferences and map-based indicators of landscape structure. *Landscape and Urban Planning* 78, 465–474.
- Dumont B, Fortun-Lamothe L, Jouven M, Thomas M and Tichit M 2013. Prospects from agroecology and industrial ecology for animal production in the 21st century. *Animal* 7, 1028–1043.
- Durant D, Tichit M, Fritz H and Kernéis E 2008. Field occupancy by breeding lapwings *Vanellus vanellus* and redshanks *Tringa totanus* in agricultural wet grasslands. *Agriculture, Ecosystems and Environment* 128, 146–150.
- European Commission 2011a. The CAP towards 2020 – impact assessment of alternative policy options. European Commission. Directorate-General for Agriculture and Rural Development, Brussels, Belgium.
- European Commission 2011b. Our life insurance, our natural capital: an EU biodiversity strategy to 2020 COM(2011) 244 final. European Commission, Brussels, Belgium.
- Foley JA, DeFries R, Asner GP, Barford C, Bonan G, Carpenter SR, Chapin FS, Coe MT, Daily GC, Gibbs HK, Helkowski JH, Holloway T, Howard EA, Kucharik CJ, Monfreda C, Patz JA, Prentice C, Ramankutty N and Snyder PK 2005. Global consequences of land use. *Science* 309, 570–574.
- Ford H, Garbutt A, Jones DL and Jones L 2012. Impacts of grazing abandonment on ecosystem service provision: coastal grassland as a model system. *Agriculture, Ecosystems and Environment* 162, 108–115.
- Franzén M and Nilsson SG 2008. How can we preserve and restore species richness of pollinating insects on agricultural land? *Ecography* 31, 698–708.
- García-Llorente M, Martín-López B, Iniesta-Arandia I, López-Santiago CA, Aguilera PA and Montes C 2012. The role of multi-functionality in social preferences toward semi-arid rural landscapes: an ecosystem service approach. *Environmental Science and Policy* 19–20, 136–146.
- Gómez-Baggethun E and de Groot R 2008. Capital natural y funciones de los ecosistemas: explorando las bases ecológicas de la economía. *Ecosistemas* 16, 1–11.
- Gómez-Baggethun E and Ruiz-Pérez M 2011. Economic valuation and the commodification of ecosystem services. *Progress in Physical Geography* 35, 613–628.
- Groot JCJ, Jellema A and Rossing WAH 2010. Designing a hedgerow network in a multifunctional agricultural landscape: balancing trade-offs among ecological quality, landscape character and implementation costs. *European Journal of Agronomy* 32, 112–119.
- Groot JCJ, Oomen GJM and Rossing WAH 2012. Multi-objective optimization and design of farming systems. *Agricultural Systems* 110, 63–77.
- Groot JCJ, Rossing WAH, Jellema A, Stobbelaar DJ, Renting H and Van Ittersum MK 2007. Exploring multi-scale trade-offs between nature conservation, agricultural profits and landscape quality-A methodology to support discussions on land-use perspectives. *Agriculture, Ecosystems and Environment* 120, 58–69.
- Groot JCJ, Rossing WAH, Tichit M, Turpin N, Jellema A, Baudry J, Verburg PH, Doyen L and van de Ven GWJ 2009. On the contribution of modelling to multi-functional agriculture: learning from comparisons. *Journal of Environmental Management* 90, S147–S160.
- Haslem A and Bennett AF 2008. Birds in agricultural mosaics: the influence of landscape pattern and countryside heterogeneity. *Ecological Applications* 18, 185–196.

- Heikkinen H, Sarkki S and Nuttall M 2012. Users or producers of ecosystem services? A scenario exercise for integrating conservation and reindeer herding in northeast Finland. *Pastoralism* 2, 1–24.
- Hicks CC, McClanahan TR, Cinner JE and Hills JM 2009. Trade-offs in values assigned to ecological goods and services associated with different coral reef management strategies. *Ecology and Society* 14(1), 10.
- Hoiss B, Gaviria J, Leingärtner A, Krauss J and Steffan-Dewenter I 2013. Combined effects of climate and management on plant diversity and pollination type in alpine grasslands. *Diversity and Distributions* 19, 386–395.
- Jouven M and Baumont R 2008. Simulating grassland utilization in beef suckler systems to investigate the trade-offs between production and floristic diversity. *Agricultural Systems* 96, 260–272.
- Kareiva P, Watts S, McDonald R and Boucher T 2007. Domesticated nature: shaping landscapes and ecosystems for human welfare. *Science* 316, 1866–1869.
- Kassie GT, Abdulai A and Wolltyn C 2009. Valuing traits of indigenous cows in central Ethiopia. *Journal of Agricultural Economics* 60, 386–401.
- Kumar P 2010. *The Economics of Ecosystems and Biodiversity: Ecological and Economic Foundations*. TEEB, Earthscan, London, UK and Washington, USA.
- Lamarque P, Tappeiner U, Turner C, Steinbacher M, Bardgett RD, Szukics U, Schermer M and Lavorel S 2011. Stakeholder perceptions of grassland ecosystem services in relation to knowledge on soil fertility and biodiversity. *Regional Environmental Change* 11, 791–804.
- Lorencová E, Frélichová J, Nelson E and Vačkář D 2013. Past and future impacts of land use and climate change on agricultural ecosystem services in the Czech Republic. *Land Use Policy* 33, 183–194.
- Ma S and Swinton SM 2011. Valuation of ecosystem services from rural landscapes using agricultural land prices. *Ecological Economics* 70, 1649–1659.
- MacDonald D, Crabtree JR, Wiesinger G, Dax T, Stamou N, Fleury P, Lazpita JG and Gibon A 2000. Agricultural abandonment in mountain areas of Europe: environmental consequences and policy response. *Journal of Environmental Management* 59, 47–69.
- Marriott CA, Fisher JM, Hood K and Pakeman RJ 2010. Impacts of extensive grazing and abandonment on grassland soils and productivity. *Agriculture, Ecosystems and Environment* 139, 476–482.
- Martin-Collado D 2013. Integration of socioeconomic and genetic aspects involved in the conservation of animal genetic resources. Thesis PhD, Universidad Politécnica de Madrid, Madrid, Spain.
- Martin G, Cruz P, Theau JP, Jouany C, Fleury P, Granger S, Faivre R, Balent G, Lavorel S and Duru M 2009. A multi-site study to classify semi-natural grassland types. *Agriculture Ecosystems and Environment* 129, 508–515.
- Martín-López B, García-Llorente M, Palomo I and Montes C 2011. The conservation against development paradigm in protected areas: valuation of ecosystem services in the Doñana social-ecological system (southwestern Spain). *Ecological Economics* 70, 1481–1491.
- Martín-López B, Gómez-Baggethun E, García-Llorente M and Montes C 2014. Trade-offs across value-domains in ecosystem services assessment. *Ecological Indicators* 37, 220–228.
- Martín-López B, Iniesta-Arandia I, García-Llorente M, Palomo I, Casado-Arzuaga I, Amo DGD, Gómez-Baggethun E, Oteros-Rozas E, Palacios-Agundez I, Willaarts B, González JA, Santos-Martín F, Onaindia M, López-Santiago C and Montes C 2012. Uncovering ecosystem service bundles through social preferences. *PLoS One* 7, e38970.
- Martinsen V, Mulder J, Austrheim G and Myrsterud A 2011. Carbon storage in low-alpine grassland soils: effects of different grazing intensities of sheep. *European Journal of Soil Science* 62, 822–833.
- McMahon BJ, Helden A, Anderson A, Sheridan H, Kinsella A and Purvis G 2010. Interactions between livestock systems and biodiversity in South-East Ireland. *Agriculture, Ecosystems and Environment* 139, 232–238.
- Medina-Roldán E, Paz-Ferreiro J and Bardgett RD 2012. Grazing exclusion affects soil and plant communities, but has no impact on soil carbon storage in an upland grassland. *Agriculture, Ecosystems and Environment* 149, 118–123.
- Michaud A, Plantureux S, Amiaud B, Carrère P, Cruz P, Duru M, Dury B, Farruggia A, Fiorelli JL, Kerneis E and Baumont R 2012. Identification of the environmental factors which drive the botanical and functional composition of permanent grasslands. *Journal of Agricultural Science* 150, 219–236.
- Millennium Ecosystem Assessment 2005. *Ecosystems and human well-being: synthesis*. Island Press, World Resources Institute, Washington, DC, USA.
- Nieto-Romero M, Oteros-Rozas E, González JA and Martín-López B 2014. Exploring the knowledge landscape of ecosystem services assessments in Mediterranean agroecosystems: insights for future research. *Environmental Science and Policy* 37, 121–133.
- Oteros-Rozas E 2013. Análisis de una práctica agraria tradicional en la cuenca mediterránea desde la perspectiva socio-ecológica: la trashumancia en la Cañada Real Conquense. Thesis PhD, Universidad Autónoma de Madrid, Madrid, Spain.
- Oteros-Rozas E, González JA, Martín-López B, López CA, Zorrilla-Miras P and Montes C 2012. Evaluating ecosystem services in transhumance cultural landscapes. An interdisciplinary and participatory framework. *GAIA* 21, 185–193.
- Oteros-Rozas E, Martín-López B, González JA, Plieninger T, López CA and Montes C 2013a. Socio-cultural valuation of ecosystem services in a transhumance social-ecological network. *Regional Environmental Change*, doi:10.1007/s10113-013-0571-y.
- Oteros-Rozas E, Martín-López B, López CA, Palomo I and González JA 2013b. Envisioning the future of transhumant pastoralism through participatory scenario planning: a case study in Spain. *The Rangeland Journal* 35, 251–272.
- Oxbrough AG, Gittings T, O'Halloran J, Giller PS and Kelly TC 2006. The initial effects of afforestation on the ground-dwelling spider fauna of Irish peatlands and grasslands. *Forest Ecology and Management* 237, 478–491.
- Palomo I, Martín-López B, Potschin M, Haines-Young R and Montes C 2013. National parks, buffer zones and surrounding lands: mapping ecosystem service flows. *Ecosystem Services* 4, 104–116.
- Pereira E, Queiroz C, Pereira HM and Vicente L 2005. Ecosystem services and human well-being: a participatory study in a mountain community in Portugal. *Ecology and Society* 10(2), 14.
- Plieninger T, Höchtl F and Spek T 2006. Traditional land-use and nature conservation in European rural landscapes. *Environmental Science and Policy* 9, 317–321.
- Plieninger T, Dijks S, Oteros-Rozas E and Bieling C 2013. Assessing, mapping, and quantifying cultural ecosystem services at community level. *Land Use Policy* 33, 118–129.
- Polasky S, Nelson E, Lonsdorf E, Fackler P and Starfield A 2005. Conserving species in a working landscape: land use with biological and economic objectives. *Ecological Applications* 15, 1387–1401.
- Pouta E and Ovaskainen V 2006. Assessing the recreational demand for agricultural land in Finland. *Agricultural and Food Science* 15, 375–387.
- Power AG 2010. Ecosystem services and agriculture: tradeoffs and synergies. *Philosophical Transactions of the Royal Society B: Biological Sciences* 365, 2959–2971.
- Raudsepp-Hearne C, Peterson GD and Bennett EM 2010. Ecosystem service bundles for analyzing tradeoffs in diverse landscapes. *Proceedings of the National Academy of Sciences of the United States of America* 107, 5242–5247.
- Ready RC and Abdalla CW 2005. The amenity and disamenity impacts of agriculture: Estimates from a hedonic pricing model. *American Journal of Agricultural Economics* 87, 314–326.
- Riedel JL, Bernues A and Casasus I 2013. Livestock grazing impacts on herbage and shrub dynamics in a Mediterranean natural park. *Rangeland Ecology and Management* 66, 224–233.
- Ruiz-Mirazo J and Robles AB 2012. Impact of targeted sheep grazing on herbage and holm oak saplings in a silvopastoral wildfire prevention system in south-eastern Spain. *Agroforestry Systems* 86, 477–491.
- Ruto E, Garrod G and Scarpa R 2008. Valuing animal genetic resources: A choice modelling application to indigenous cattle in Kenya. *Agricultural Economics* 38, 89–98.
- Sabatier R, Doyen L and Tichit M 2010. Modelling trade-offs between livestock grazing and wader conservation in a grassland agroecosystem. *Ecological Modelling* 221, 1292–1300.
- Sabatier R, Doyen L and Tichit M 2012. Action versus result-oriented schemes in a grassland agroecosystem: a dynamic modelling approach. *PLoS One* 7, e33257.
- Sabatier R, Doyen L and Tichit M 2013. Heterogeneity and the trade-off between ecological and productive functions of agro-landscapes: a model of cattle-bird interactions in a grassland agroecosystem. *Agricultural Systems*, doi:10.1016/j.agsy.2013.02.008. Published online 25 March 2013.

- Scarpa R, Ruto ESK, Kristjanson P, Radeny M, Drucker AG and Rege JEO 2003. Valuing indigenous cattle breeds in Kenya: An empirical comparison of stated and revealed preference value estimates. *Ecological Economics* 45, 409–426.
- Scoones I 1992. The economic value of livestock in the communal areas of southern Zimbabwe. *Agricultural Systems* 39, 339–359.
- Seppelt R, Dormann CF, Eppink FV, Lautenbach S and Schmidt S 2011. A quantitative review of ecosystem service studies: approaches, shortcomings and the road ahead. *Journal of Applied Ecology* 48, 630–636.
- Shonkwiler JS and Englin J 2005. Welfare losses due to livestock grazing on public lands: a count data systemwide treatment. *American Journal of Agricultural Economics* 87, 302–313.
- Souchère V, Millair L, Echeverria J, Bousquet F, Le Page C and Etienne M 2010. Co-constructing with stakeholders a role-playing game to initiate collective management of erosive runoff risks at the watershed scale. *Environmental Modelling and Software* 25, 1359–1370.
- Soussana JF, Loiseau P, Vuichard N, Ceschia E, Balesdent J, Chevallier T and Arrouays D 2004. Carbon cycling and sequestration opportunities in temperate grasslands. *Soil Use and Management* 20, 219–230.
- Stoate C, Boatman ND, Borralho RJ, Carvalho CR, De Snoo GR and Eden P 2001. Ecological impacts of arable intensification in Europe. *Journal of Environmental Management* 63, 337–365.
- Struhsaker TT, Struhsaker PJ and Siex KS 2005. Conserving Africa's rain forests: problems in protected areas and possible solutions. *Biological Conservation* 123, 45–54.
- Swinton SM, Hamilton SK, Lupi F and Robertson GP 2007. Special issue: ecosystem services and agriculture. *Ecological Economics* 64, 245–385.
- Tasser E, Walde J, Tappeiner U, Teutsch A and Noggler W 2007. Land-use changes and natural reforestation in the Eastern Central Alps. *Agriculture, Ecosystems and Environment* 118, 115–129.
- Tichit M, Durant D and Kernéis E 2005a. The role of grazing in creating suitable sward structures for breeding waders in agricultural landscapes. *Livestock Production Science* 96, 119–128.
- Tichit M, Renault O and Potter T 2005b. Grazing regime as a tool to assess positive side effects of livestock farming systems on wading birds. *Livestock Production Science* 96, 109–117.
- Tichit M, Puillet L, Sabatier R and Teillard F 2011. Multicriteria performance and sustainability in livestock farming systems: functional diversity matters. *Livestock Science* 139, 161–171.
- Tichit M, Doyen L, Lemel JY, Renault O and Durant D 2007. A co-viability model of grazing and bird community management in farmland. *Ecological Modelling* 206, 277–293.
- Varah A, Jones H, Smith J and Potts SG 2013. Enhanced biodiversity and pollination in UK agroforestry systems. *Journal of the Science of Food and Agriculture* 93, 2073–2075.
- Venkatachalam L 2004. The contingent valuation method: a review. *Environmental Impact Assessment Review* 24, 89–124.
- Weigelt A, Weisser WW, Buchmann N and Scherer-Lorenzen M 2009. Biodiversity for multifunctional grasslands: equal productivity in high-diversity low-input and low-diversity high-input systems. *Biogeosciences* 6, 1695–1706.
- Zander KK and Drucker AG 2008. Conserving what's important: Using choice model scenarios to value local cattle breeds in East Africa. *Ecological Economics* 68, 34–45.
- Zander KK, Signorello G, De Salvo M, Gandini G and Drucker AG 2013. Assessing the total economic value of threatened livestock breeds in Italy: implications for conservation policy. *Ecological Economics* 93, 219–229.
- Zhang W, Ricketts TH, Kremen C, Carney K and Swinton SM 2007. Ecosystem services and dis-services to agriculture. *Ecological Economics* 64, 253–260.